

Review

- Knowledge Base
- Propositional Logic
- First Order Logic
 - Universal and Existential Quantifier
- Reduction of first-order inference to propositional inference
 - Universal and Existential Instantiation

Probability

Uncertainty and Vagueness

- Real world is not always exact and certain
 - Do the best with what is already known
 - Agent needs to draw conclusions when available information is uncertain, vague, or incomplete

Uncertainty and Vagueness

- Real world is not always exact and certain
 - Do the best with what is already known
 - Agent needs to draw conclusions when available information is uncertain, vague, or incomplete
- Sources of uncertainty
 - Unreliable data
 - Defective measurement device
 - Incomplete data
 - Only partial data available
 - Imprecise data/rules
 - Approximations of data
 - Rules for drawing conclusions may also be imprecise

Handling Uncertain Knowledge

- Consider dental diagnosis using first-order logic

$$\forall p \text{ Symptom}(p, \text{Toothache}) \rightarrow \text{Disease}(p, \text{Cavity})$$

- Rule is wrong!
 - Not all patients with toothache have cavities

Handling Uncertain Knowledge

- Consider dental diagnosis using first-order logic

$$\forall p \text{ Symptom}(p, \text{Toothache}) \rightarrow \text{Disease}(p, \text{Cavity})$$

- Rule is wrong!
 - Not all patients with toothache have cavities
- Need to add an almost unlimited list of possible causes

$$\begin{aligned} \forall p \text{ Symptom}(p, \text{Toothache}) \rightarrow & \text{Disease}(p, \text{Cavity}) \\ & \vee \text{Disease}(p, \text{GumDisease}) \\ & \vee \text{Disease}(p, \text{ImpactedWisdom}) \dots \end{aligned}$$

Handling Uncertain Knowledge

- Use of first-order logic fails in diagnosis for three reasons
 - Laziness
 - Theoretical ignorance
 - Practical ignorance

Handling Uncertain Knowledge

- Use of first-order logic fails in diagnosis for three reasons
 - Laziness
 - Too much work to list all possible consequents
 - Theoretical ignorance
 - Practical ignorance

Handling Uncertain Knowledge

- Use of first-order logic fails in diagnosis for three reasons
 - Laziness
 - Too much work to list all possible consequents
 - Theoretical ignorance
 - No complete theory for domain available
 - Practical ignorance

Handling Uncertain Knowledge

- Use of first-order logic fails in diagnosis for three reasons
 - Laziness
 - Too much work to list all possible consequents
 - Theoretical ignorance
 - No complete theory for domain available
 - Practical ignorance
 - Even if have all rules, may be uncertain about a patient (all tests have not been run to get data)

Handling Uncertain Knowledge

- System's knowledge (Knowledge Base) can **only** provide a degree of belief about sentences
 - Sentences are in fact either true or false

Handling Uncertain Knowledge

- System's knowledge (Knowledge Base) can **only** provide a degree of belief about sentences
 - Sentences are in fact either true or false
- Probability theory
 - Deals with uncertainty
 - Assignment of probability depends on percepts received to date (evidence)
 - Assigns numerical degree of belief between 0-1

Handling Uncertain Knowledge

- System's knowledge (Knowledge Base) can **only** provide a degree of belief about sentences
 - Sentences are in fact either true or false
- Probability theory
 - Deals with uncertainty
 - Assignment of probability depends on percepts received to date (evidence)
 - Assigns numerical degree of belief between 0-1
 - Probability of 0 absolute belief that sentence is FALSE
 - Probability of 1 is absolute belief that sentence is TRUE
 - Probability of 0.8 (80%) that patient has cavity if has toothache

Basics of Probability

- Prior probability
- Conditional probability
- Bayes rule

Basic Concepts of Probability

- Prior probability
 - *A priori* initial information: $P(A)$
 - “Probability of event A occurring”

Basic Concepts of Probability

- Prior probability
 - *A priori* initial information: $P(A)$
 - “Probability of event A occurring”
 - Independent of any evidence

Basic Concepts of Probability

- Joint probability: $P(A \wedge B)$
 - “Probability of event A and B occurring together”
 - Also often written $P(A, B)$

Basic Concepts of Probability

- Conditional probability
 - Information based on additional evidence: $P(A | C)$
 - “Probability of A, given that I know C”

Prior Probability

Prior Probability

- Probability assessment before any evidence obtained
 - “Probability of event A ” $\rightarrow P(A)$
 - e.g., $P(Cavity) = 0.1$
 - In the absence of any other information, 10% chance of the patient having a cavity

Prior Probability

- Probability assessment before any evidence obtained
 - “Probability of event A” $\rightarrow P(A)$
 - e.g., $P(\text{Cavity}) = 0.1$
 - In the absence of any other information, 10% chance of the patient having a cavity

- Random variable assignments of exclusive values

$$\left. \begin{array}{l} P(\text{weather} = \text{Sunny}) = 0.7 \\ P(\text{weather} = \text{Rainy}) = 0.2 \\ P(\text{weather} = \text{Cloudy}) = 0.08 \\ P(\text{weather} = \text{Snow}) = 0.02 \end{array} \right\} \text{ Must add up to 1}$$

$\text{weather} \rightarrow \{\text{Sunny}, \text{Rainy}, \text{Cloudy}, \text{Snow}\}$

Prior Probability

- Properties defining probability $P(A)$

$$0 \leq P(A) \leq 1$$

$$P(\text{"certain event"}) = 1$$

$$P(A) + P(\neg A) = 1$$

$$P(\neg A) = 1 - P(A)$$

Prior Probability

- Properties defining probability $P(A)$

$$0 \leq P(A) \leq 1$$

$$P(\text{"certain event"}) = 1$$

$$P(A) + P(\neg A) = 1$$

$$P(\neg A) = 1 - P(A)$$

- Simple die examples

- Each side of die labeled from 1 to 6

- Let event A be a die stops with 1 showing on top: $P(A) = 1/6$

- Let event $\sim A$ be a die stops with numbers other than a 1 showing on top

$$P(\neg A) = ?$$

$$P(\neg A) = 1 - P(A)$$

$$= 1 - 1/6$$

$$= 5/6$$

Joint Probability

- Joint probability (“and”)
 $P(Cavity \wedge \neg Insured) = 0.06$

Prior & Joint Probability

- Joint probability (“and”)


$$P(Cavity \wedge \neg Insured) = 0.06$$

- Marginalization

$$P(Cavity) = P(Cavity \wedge Insured) + P(Cavity \wedge \neg Insured)$$

Probability axiom of a Disjunction (“or”)

$$P(A \vee B) = P(A) + P(B) - P(A \wedge B)$$



Conditional Probability

Conditional Probability

- Probability assessment after evidence is obtained (“posterior probability”)
 - “Probability of event A given we know B”
 $\rightarrow P(A \mid B)$
 - e.g., $P(\text{Cavity} \mid \text{Toothache}) = 0.8$
 - If observe patient with toothache, 80% chance patient has cavity

Conditional Probability

- Probability assessment after evidence is obtained (“posterior probability”)
 - “Probability of event A given we know B”
 $\rightarrow P(A | B)$
 - e.g., $P(\text{Cavity} | \text{Toothache}) = 0.8$
 - If observe patient with toothache, 80% chance patient has cavity
- Properties defining conditional probability $P(A|B)$

$$0 \leq P(A|B) \leq 1$$

$$P(A|B) + P(\neg A|B) = 1$$

$$P(\neg A|B) = 1 - P(A|B)$$

Conditional Probability

- Probability assessment after evidence is obtained (“posterior probability”)
 - “Probability of event A given we know B”
 $\rightarrow P(A | B)$
 - e.g., $P(\text{Cavity} | \text{Toothache}) = 0.8$
 - If observe patient with toothache, 80% chance patient has cavity
- Properties defining conditional probability $P(A|B)$

$$0 \leq P(A|B) \leq 1$$

$$P(A|B) + P(\neg A|B) = 1$$

$$P(\neg A|B) = 1 - P(A|B)$$

$$P(A|\neg B) + P(\neg A|\neg B) = 1$$

$$P(\neg A|\neg B) = 1 - P(A|\neg B)$$

Conditional Probability

- Probability assessment after evidence is obtained (“posterior probability”)
 - “Probability of event A given we know B”
 $\rightarrow P(A | B)$
 - e.g., $P(\text{Cavity} | \text{Toothache}) = 0.8$
 - If observe patient with toothache, 80% chance patient has cavity
- Properties defining conditional probability $P(A|B)$

$$0 \leq P(A|B) \leq 1$$

$$P(A|B) + P(\neg A|B) = 1$$

$$P(\neg A|B) = 1 - P(A|B)$$

$$P(A|\neg B) + P(\neg A|\neg B) = 1$$

$$P(\neg A|\neg B) = 1 - P(A|\neg B)$$

No relation between $P(A|B)$ and $P(A|\neg B)$

Conditional Probability

- Conditional probability in terms of unconditionals

$$P(A \wedge B) = P(A | B) \cdot P(B) \quad [\text{Product or multiply rule}]$$

thus

$$P(A | B) = P(A \wedge B) / P(B)$$

Conditional Probability

- Conditional probability in terms of unconditionals

$$P(A \wedge B) = P(A | B) \cdot P(B) \quad [\text{Product or multiply rule}]$$

thus

$$P(A | B) = P(A \wedge B) / P(B)$$

Note: If B independent of A, changes to B do not affect A:

$$P(A | B) = P(A | \neg B) = P(A)$$

$$P(A \wedge B) = P(A | B) \cdot P(B) = P(A)P(B)$$

Conditional Probability

- As soon as get more information C , must compute new conditional

$$P(A \mid B \wedge C)$$

Conditional Probability

- As soon as get more information C , must compute new conditional
$$P(A \mid B \wedge C)$$
- If A and B are conditionally independent given evidence C
$$P(A \mid B \wedge C) = P(A \mid C)$$

Inference from Joint Probabilities

- Table of 2 propositions: *Cavity* and *Toothache*

	<i>Toothache</i>	\neg <i>Toothache</i>
<i>Cavity</i>	0.04	0.06
\neg <i>Cavity</i>	0.01	0.89

$\rightarrow P(\text{Cavity} \wedge \neg \text{Toothache})$

$P(\text{Toothache})$

$$\begin{aligned} P(\text{Cavity} \vee \text{Toothache}) &= P(\text{Cavity}) + P(\text{Toothache}) - P(\text{C} \wedge \text{T}) \\ &= (0.04 + 0.06) + (0.04 + 0.01) - 0.04 = 0.11 \end{aligned}$$

$$\begin{aligned} P(\text{Cavity} \mid \text{Toothache}) &= P(\text{Cavity} \wedge \text{Toothache}) / P(\text{Toothache}) \\ &= 0.04 / (0.04 + 0.01) = 0.80 \end{aligned}$$

Bayes Rule

Bayes Rule

- Typical problem is to evaluate a hypothesis H given evidence/data E :
 - $P(H \mid E)$

Bayes Rule

- Typical problem is to evaluate a hypothesis H given evidence/data E :
 - $P(H | E)$
- Recall our earlier product rule for a conditionals

$$P(H | E) = P(H \wedge E) / P(E)$$

But can be hard to get or understand $P(H \wedge E)$

Bayes Rule

- Typical problem is to evaluate a hypothesis H given evidence/data E :
 - $P(H | E)$
- Recall our earlier product rule for conditionals

$$P(H | E) = P(H \wedge E) / P(E)$$

But can be hard to get or understand $P(H \wedge E)$

- Recall product rule (multiplicative law)

$$P(A \wedge B) = P(A | B) \cdot P(B)$$

$$P(B \wedge A) = P(B | A) \cdot P(A)$$

Bayes Rule

- Typical problem is to evaluate a hypothesis H given evidence/data E :
 - $P(H \mid E)$
- Recall our earlier product rule for conditionals

$$P(H \mid E) = P(H \wedge E) / P(E)$$

But can be hard to get or understand $P(H \wedge E)$

- Recall product rule (multiplicative law)

$$P(A \wedge B) = P(A \mid B) \cdot P(B)$$

$$P(B \wedge A) = P(B \mid A) \cdot P(A)$$

[equate right-hand sides $[P(A \wedge B) = P(B \wedge A)]$]

Bayes Rule

- Typical problem is to evaluate a hypothesis H given evidence/data E :
 - $P(H | E)$
- Recall our earlier product rule for conditionals

$$P(H | E) = P(H \wedge E) / P(E)$$

But can be hard to get or understand $P(H \wedge E)$

- Recall product rule (multiplicative law)

$$P(A \wedge B) = P(A | B) \cdot P(B)$$

$$P(B \wedge A) = P(B | A) \cdot P(A)$$

[equate right-hand sides $[P(A \wedge B) = P(B \wedge A)]$]

$$P(A | B) \cdot P(B) = P(B | A) \cdot P(A)$$

$$P(A | B) = P(B | A) \cdot P(A) / P(B)$$

$$P(H | E) = P(E | H) \cdot P(H) / P(E) \longleftarrow \text{Bayes' rule}$$

Bayes Rule

- Other variations of denominator

$$\text{BAYES: } P(H | E) = P(E | H)P(H) / P(E)$$

$$\begin{aligned} P(E) &= P(E \wedge H) + P(E \wedge \neg H) \\ &= P(E | H)P(H) + P(E | \neg H)P(\neg H) \\ &= \sum_i P(E | H_i)P(H_i) \end{aligned}$$

Bayes Rule

- Other variations of denominator

$$\text{BAYES: } P(H | E) = P(E | H)P(H) / P(E)$$

$$\begin{aligned} P(E) &= P(E \wedge H) + P(E \wedge \neg H) \\ &= P(E | H)P(H) + P(E | \neg H)P(\neg H) \\ &= \sum_i P(E | H_i)P(H_i) \end{aligned}$$

“sum/integrate over all possible hypotheses”

Bayes Rule

- Other variations of denominator

$$\text{BAYES: } P(H | E) = P(E | H)P(H) / P(E)$$

$$\begin{aligned} P(E) &= P(E \wedge H) + P(E \wedge \neg H) \\ &= P(E | H)P(H) + P(E | \neg H)P(\neg H) \\ &= \sum_i P(E | H_i)P(H_i) \end{aligned}$$

“sum/integrate over all possible hypotheses”

$$P(H | E) = \frac{P(E | H)P(H)}{\sum_i P(E | H_i)P(H_i)}$$

Summary

- Decisions not always exact and certain
 - Agent needs to draw conclusions when available information is uncertain
- Probability laws
 - Prior probability
 - A priori initial information, independent of experience
 - Joint probability
 - Conditional probability
 - Information based on additional evidence
 - Bayes rule
 - Updates belief measures in response to evidence