Apr 02, 2019

Working

E gene amplification

Bo Yi¹, Yi Chen¹, Xiao Ma¹, Haibin Wang², Rong Wang¹, Keqin Ding¹, Lei Xie¹, Dongliang Zhang¹, Shuli Jiao¹, Xuying Lao¹, Yi-Chen Chiang³, Yanhua Su³, Benhua Zhao³, Guozhang Xu¹, Tianmu Chen³

¹Ningbo Municipal Center for Disease Control and Prevention, ²Haishu District Center for Disease Control and Prevention, ³State Key Laboratory of Molecular Vaccinology and Molecular Diagnostics, School of Public Health, Xiamen University

dx.doi.org/10.17504/protocols.io.zpuf5nw



Tianmu Chen



ABSTRACT

PROTOCOL STATUS

Working

We use this protocol in our group and it is working

- The dengue virus strains obtained from cell culture were extracted by TGuide S32 magnetic bead method DNA/RNA extraction kit of Tiangen Biochemical Technology (Beijing) Co., Ltd. and amplified by one step RNA PCR Kit (Code No: DRR057A) reagent of Bao Bioengineering (Dalian) Co., Ltd.
- The full sequence of E gene amplified by primer sequence was D1 (E) F: CAA GAA CCG AAA CA / GT GGA TGT C; D1 (E) R: GGC TGA TCG AAT TCC ACA CAC, and the length of amplified fragment was 1849 bp.
- Reaction conditions: reverse transcription at 50 °C 30 minutes, reverse transcription at 94 °C 2 minutes, reverse transcription at 94 °C 30 seconds, reverse transcription at 52°C 30 seconds, reverse transcription at 72°C 2 minutes, 40 cycles and extension at 72°C. The amplified product was 5 μL, and 1.5% agarose gel electrophoresis was used to confirm the reaction product according to Marker position.

This is an open access protocol distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited