



May 15,  
2019

Working

## UC Davis - LDL Protocol [↗](#)

Peter Havel<sup>1</sup>

<sup>1</sup>University of California, Davis

dx.doi.org/10.17504/protocols.io.yrcfv2w

Mouse Metabolic Phenotyping Centers  
Tech. support email: [info@mmpc.org](mailto:info@mmpc.org)

Lili Liang

### ABSTRACT

#### Summary:

LDL and VLDL are separated from HDL using a precipitation reagent. Then the HDL fraction is measured for either TC or TG using the same reagents for total cholesterol or triglyceride.

### EXTERNAL LINK

<https://mmpc.org/shared/document.aspx?id=94&docType=Protocol>

### MATERIALS

NAME	CATALOG #	VENDOR
Calibrator	TR43002	Fisher Diagnostics
TC Reagents	TR13421	Fisher Diagnostics
TG Reagents	TR22421	Fisher Diagnostics
2X LDL/VLDL Precipitation Buffer	ab105138	Abcam
PBS		
Microplate		
Platereader		

### MATERIALS TEXT

#### Note:

Fisher Scientific, [RRID:SCR\\_008452](#)

Abcam, [RRID:SCR\\_012931](#)

- 1 Add 25µl 2X precipitation buffer to 25µl of sample using a positive displacement pipet.
- 2 Vortex and let sit at RT for 10 minutes.
- 3 Centrifuge at 2000×g for 10 minutes at 4°C.

4 Pipet supernatant into new tube, this is the HDL fraction.

5 Add 5 µl of calibrator and sample to each well.

**IMPORTANT: Make sure not to add any bubbles to the wells when dispensing reagents, this will interfere with reading in the platereader.**

6 Add 300 µl of TC or TG reagent to each well. Incubate at 37°C for 5 minutes. Read at 540 nm.

**IMPORTANT: If samples are hemolyzed, pipet a blank well with 5µl sample and 300µl PBS**

7 Subtract blank readings from final readings. The assay will be linear so the unknown samples can be calculated as (Sample Absorbance ÷ Calibrator Absorbance) × Calibrator Concentration.

8 HDL samples are diluted ½ so multiply these by 2 to get the final value. Subtract this from the total triglyceride or cholesterol value to get the LDL/VLDL value.



This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited