



Aug 14,
2019

Biochemical Measures of Neuropathy - Genotyping [↗](#)

Eva Feldman¹

¹University of Michigan - Ann Arbor

1 Works for me [dx.doi.org/10.17504/protocols.io.3qdgms6](https://doi.org/10.17504/protocols.io.3qdgms6)

Diabetic Complications Consortium

Tech. support email: rmcindoe@augusta.edu



Lili Liang

ABSTRACT

Summary:

Oxidative stress is highly correlated with the metabolic changes caused by hyperglycemia. Increased levels of glucose overload mitochondria and result in the production of reactive oxygen species (ROS). In addition, the flow of excess glucose through cellular pathways decreases the cell's normal ability to detoxify ROS. As a result, the neurons and axons of the peripheral nervous system contain increased levels of ROS and decreased antioxidant capacity. The following assays are used to measure these changes in rodent models of diabetic neuropathy.

Diabetic Complication:



Neuropathy

EXTERNAL LINK

<https://www.diacomp.org/shared/document.aspx?id=54&docType=Protocol>

MATERIALS

NAME	CATALOG #	VENDOR
10X PCR Buffer	10966-034	Life Technologies
50 mM MgCl ₂	10966-034	Life Technologies
10 mM dNTPs	10297-018	Life Technologies
Taq DNA Polymerase (5 U/μL)		Life Technologies
0.5 mL thin-walled tubes	LS-9350-X	Life Science Products
Forward primer (10 μM)		
Reverse primer (10 μM)		
DNA Template (200 ng/μL)		
Sterile deionized H ₂ O		

MATERIALS TEXT

Note:

Life Technologies ([RRID:SCR_008817](#))

PCR Amplification:

- 1 Prepare a Master Mix containing the following volumes of reagents:

Reagent	μL
10X PCR Buffer	2.5
50 mM MgCl_2	1.25
dNTPs	2.5
M13 Forward Primer	1.0
M13 Reverse Primer	1.0
<i>Taq</i> DNA Polymerase	0.2
DNA Template	2.0
Sterile dd-H ₂ O	14.55
Total Volume =	25.00

- 2 Into each 0.5mL thin-walled tube, pipette 23.0- μL of Master Mix.
- 3 Pipet 2.0 μL of DNA template (200 ng/ μL) into thin-walled tube. Mix gently, and spin down.
- 4 Amplify using the following program:

94°C, 5 min. → 94°C, 1 min. → 64°C, 1 min. → 72°C, 1.5 min. → 72°C, 10 min. → 4°C, soak
Repeat 35 times

- 5 Verify PCR product by separation on 2% agarose.
- 6 Store tubes at -20°C until needed. If plates will be used within a few days they may be stored at 4°C.



This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited