



Preparation of electrocompetent cells

joosef Hoff¹

¹Phillips University

dx.doi.org/10.17504/protocols.io.u3heyj6

iGEM Marburg 2018

joosef Hoff ⚡

ABSTRACT

Preparation of electrocompetent cells

Weinstock paper:

[Matthew T Weinstock](#), [Eric D Heseke](#), [Christopher M Wilson](#), [Daniel G Gibson](#)

Vibrio natriegens as a fast-growing host for molecular biology

Nature Methods volume 13, pages 849–851 (2016)

To prepare the day before:

- LB I media +V2 salts (204 mM NaCl, 4.2 mM KCl, 23.14mM MgCl₂)
- electroporation buffer (680 mM sucrose, 7 mM K₂HPO₄, pH 7) (sterile filtrated)
- over night culture of V.n from the cryo-stock in LBI + v2 salts (37 °C ; at 200 r.p.m)

Preparing culture :

Depending on how many aliquds you want to have, incubate media with your overnight culture for a starting OD of 0.05.

The culture is grown at 37 °C in a baffled flask, shaking at 200 r.p.m. until an OD₆₀₀ between 0.5 to 0.8 is reached.

*be careful when they reach an OD near 0.1, *V. natriegens* very fast growing, so start measuring in shorter time periods.

Prechill the electroporation buffer

Washing:

From here on try always to keep your culture on ice.

- The culture is then put on ice for 15 min.
(the original Protocol sais to directly fill them into your prechilled centrifugation containments)
- The cells are pelleted at 3000x g. for 20 min at 4 °C.
- The supernatant is carefully decanted and the cell pellets are gently suspended in 10mL of **chilled** electroporation buffer.
- The suspensions are transferred to a **chilled** 50mL falcon tube and the tube is filled top with additional **chilled** electroporation buffer (50mL) and inverted several times.
- The cells are centrifuged down at 3000x g for 15 min at 4 °C.
- The wash is repeated two times for a total of three washes.

Aliquotation:

After the final wash, the supernatant is carefully decanted the cells are gently resuspended in residual electroporation buffer.

Measure the OD in a 1/20 dilution against electroporation buffer.

The volume is adjusted with additional electroporation buffer to bring the final OD₆₀₀ to 16.

The Cells are aliquoted (80µL) into chilled 1.5µL centrifugation tubes, directly frozen in liquid nitrogen and stored at -80 °C until use.

T AGS

Vibrio

electroporation

Show tags



Preparation of
electrocompetent
cells.pdf

PROTOCOL STATUS

Working

We use this protocol in our group and it is working



This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited