



## Preparing 10 L of M9 buffer for nematode culture V.3

Gurdon Institute media kitchen<sup>1</sup>

<sup>1</sup>Cancer Research UK / Wellcome Gurdon Institute



dx.doi.org/10.17504/protocols.io.5wjg7cn



## ABSTRACT

Prepare 10 litres of M9 solution for collecting and washing C. elegans.

MATERIALS

| NAME ~  | CATALOG # | VENDOR ~          |
|---|-----------|-------------------|
| double distilled water (ddH20)  |           |                   |
| Di-Sodium Hydrogen Orthophosphate Dihydrate Certified AR for Analysis Fisher Chemical | S/4450/53 |                   |
| Sodium chloride meets analytical specification of Ph.Eur Fisher Chemical              | S/3160/65 | Fisher Scientific |
| Magnesium Sulfate Heptahydrate Certified AR for Analysis Fisher Chemical              | M/1050/53 | Fisher Scientific |
| Potassium Dihydrogen Orthophosphate Certified AR for Analysis Fisher Chemical         | P/4800/53 | Fisher Scientific |
| SYCHEM autoclave  | View      | Syschem           |

## SAFETY WARNINGS

This protocol makes use of an autoclave. Make sure you know how to use it properly before starting the protocol.

1

| Ingredients                                  | Quantity  |     |
|--|-----------|-----|
| Na2HPO4                                      |           | 60g |
| KH2PO4                                       |           | 30g |
| NaCl   |           | 50g |
| Double distilled H2O                         | up to 10L |     |
|  |           |     |
| Add 1M MgSO4 after autoclaving @ 100ul/100ml |           |     |

M9\_buffer\_10L.xls

2

| 1 | Dissolve ingredients in 8L double distilled H2O                                   |
|---|---|
| 2 | Adjust volume to 10L using double distilled H2O                                   |
| 3 | Measure or use pump to dispense aliquots accurately.                              |
| 4 | Label, date and autoclave.  |
| 5 | After autoclaving, add 100ul 1M MgSO4 per 100ml. (ie. 100ul to 100ml and 200ul to |
|   | 200ml bottles).   |

protocols.io

15m

20m

| NB | 1M MgSO4 at 1ul/1ml in                           |
|----|--|
|    | the M9 buffer gives a final concentration of 1mM |

The autoclave is SYCHEM VS Series. The media cycle is 15 minutes at 121 degrees centigrade. It is then cooled down in the machine for 3 hours.

10m

After autoclaving, add 100  $\mu$ l 1M MgSO4 per 100 ml. (ie. 100  $\mu$ l to 100 ml and 200  $\mu$ l to 200 ml bottles).

NB: 1M MgSO4 at 1  $\mu$ l/1ml in the M9 buffer gives a final concentration of 1 mM

This is an open access protocol distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited