



## ChIP protocol

Krishnaprasad G. Nandanan<sup>1</sup>, Ajith V. Pankajam<sup>1</sup>, Miki Shinohara<sup>2</sup>, Parijat Chakraborty<sup>1</sup>, Gen Lin<sup>3</sup>, Sagar Salim<sup>1</sup>, Lars M. Steinmetz<sup>4</sup>, Akira Shinohara<sup>5</sup>, Koodali T. Nishant<sup>6</sup>

<sup>1</sup>School of Biology, Indian Institute of Science Education and Research Thiruvananthapuram, Trivandrum 695016, India, <sup>2</sup>Department of Advanced Bioscience, Kindai University, Nara 631-8505, Japan, <sup>3</sup>Genome Biology Unit, European Molecular Biology Laboratory, Heidelberg 69117, <sup>4</sup>Genome Biology Unit, European Molecular Biology Laboratory, Heidelberg 69117, Germany Department of Genetics, Stanford University, Stanford, CA 94305, USA, Stanford Genome Technology Center, Palo Alto, CA 94304, USA, <sup>5</sup>Institute for Protein Research, Osaka University, Osaka 565-0871, Japan, <sup>6</sup>School of Biology and Center for Computation, Modelling and Simulation, Indian Institute of Science Education and Research Thiruvananthapuram, Trivandrum 695016, India

Sep 14, 2018 dx.doi.org/10.17504/protocols.io.tkyekxw



Krishnaprasad Gurukripa Nandanan ⚡



Chromatin  
Immunoprecipitation  
method.docx

### PROTOCOL STATUS

#### Working

We use this protocol in our group and it is working



This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited