



Apr 24, 2019

Working

Transfection by Electroporation in *Euplotes crassus*

Version 2

Angela Piersanti¹¹University of Camerino[dx.doi.org/10.17504/protocols.io.2a9gah6](https://doi.org/10.17504/protocols.io.2a9gah6)

Protist Research to Optimize Tools in Genetics (PROT-G)

Angela Piersanti
University of Camerino

- 1 2x10⁴ *Euplotes crassus* cells were collected and resuspended in 0.3 M glucose solution (7% sea water and 93% of 0.3 M glucose solution).
- 2 Each round of transfection 250 µl of cells were used. 0.25 µg of Label IT® Plasmid Delivery Control Cy@3 (Mirus) were added alone or mixed with 2.5 µl of Lipofectamine® 2000 Transfection Reagent (Invitrogen) according to the supplier.
- 3 The sample was transferred to the 0.2 cm cuvette. Bio-Rad Gene Pulser was used. Conditions were set as follows: 0.2 kV, 25 µFD, 100 Ω. Time constant around 1.2.
- 4 More than 50% of cells were viable after electroporation (few cells fused together), then cells were resuspended in 3 ml of sea water.
- 5 The plasmid was visible in the cytoplasm immediately after the electroporation by fluorescent microscope.



This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited