

Mar 18,
2019

Working

Protocol for sampling and transport of nose- and thoratsamples in the Fit Futures study

Anne-Sofie Furberg¹, Karina Olsen², Gunnar Skov Simonsen², Johanna UE Sollid³, Lars Småbrekke⁴, Guri Grimnes⁵, Cristopher Sievert Nielsen⁶

¹Department of Community Medicine, Faculty of Health Sciences, UiT The Arctic University of Norway, ²Department of Microbiology and Infection Control, University Hospital of North Norway, ³Research Group for Host-Microbe Interaction, Department of Medical Biology, ⁴Department of Pharmacy, Faculty of Health Sciences, UiT The Arctic University of Norway, ⁵Endocrinology Research Group, Department of Clinical Medicine, Faculty of Health Sciences, UiT The Arctic University of Norway, ⁶Division of Ageing and Health, Norwegian Institute of Public Health

[dx.doi.org/10.17504/protocols.io.za9f2dw](https://doi.org/10.17504/protocols.io.za9f2dw)

 **Dina Stensen**
University hospital of North Norway 

ABSTRACT




The aim of the study is to evaluate prevalence of *Staphylococcus aureus* in a youth population. This samples will be connected to other possible riskfactors for *Staphylococcus aureus* carriage.

PROTOCOL STATUS

Working

We use this protocol in our group and it is working

MATERIALS

NAME 	CATALOG # 	VENDOR 
Sterile NaCl 9 mg/ml (0.9%)	View	B.Braun Medical Inc
Sterile rayon-tipped swab	View	Puritan
Transport medium - Amies Copan	View	
Tongue spatula - wood	View	
Hand disinfection	View	Antibac

MATERIALS TEXT

4 Sterile rayon-tipped swabs for each participant

SAFETY WARNINGS

BEFORE STARTING

Wash hands with soap and water or use hand disinfection to prevent contamination of the samples. The examiner should use facemask to protect the participant if necessary. If the participant has secretion from the nasal cavity, the participant should blow ones nose before sampling.

Nasal samples

- 1 Moisture the rayon-tipped swab in 2-3 drops of sterile NaCl 0,9 mg/ml. There should be two samples from the nasal cavity.
- 2 Lead the rayon-tipped swab carefully into the nasal cavity approximately 1-2 cm and rotate with delicate pressure 3 times in both nasal cavity. The same swab should be used for both anterior nares.

- 3 Place both swabs in the Coban transport medium.
- 4 Label the transport medium with testnumber and nasal swab 1 and 2.
- 5 Wash hands with soap and water or use hand disinfection after procedure

Throat samples

- 6 Moisture the rayon-tipped swab in 2-3 drops of sterile NaCl 0,9 mg/ml. There should be two samples from the throat.
- 7 Ask the participant to open his/her mouth. Press the tongue spatula against the tongue and ask the participant to say "aaa". This will simplify the sampling because this will better the view of the tonsil region.
- 8 Lead the rayon-tipped swab into the oral cavity and swab with moderate pressure on both tonsil regions (palatine tonsils). If the participants has had a former tonsillectomy/tonsillotomy the swab should be taken from the area of the removed tonsils. Use the same swab on both tonsils.
- 9 Place both swabs in the Coban transport medium.
- 10 Label the transport medium with testnumber and throat swab 1 and 2.
- 11 Wash hands with soap and water or use hand disinfection after procedure

Transport

- 12 The swabs should be placed in a suitable rack. If not picked up immediately, the samples should be placed in refrigerator awaiting pickup.



This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited