

# Small RNA interference (siRNA)( S100A6-siRNA) and cell transfection

Shanshan Feng

## Abstract

**Citation:** Shanshan Feng Small RNA interference (siRNA)( S100A6-siRNA) and cell transfection. **protocols.io**  
dx.doi.org/10.17504/protocols.io.mqfc5tn

**Published:** 18 Jan 2018

## Protocol

### Step 1.

Shortly before transfection, seed  $1 \times 10^6$  cells into each well of 6-well plates and cultured for 48 h.

### Step 2.

For the short time until transfection, incubate the cells under normal growth conditions (typically 37°C and 5% CO<sub>2</sub>).

### Step 3.

Cells were transfected with 7.5 µmol/L of each siRNA for 24 hrs in Opti-MEM medium without antibiotics and 5% fetal calf serum using Lipo-fectamine 2000 (Invitrogen, Carlsbad, USA)

### Step 4.

Incubate for 5 min at room temperature (20°C) to allow the formation of transfection complexes.

### Step 5.

Add the complexes drop-wise onto the cells. Gently swirl the plate to ensure uniform distribution of the transfection complexes.

### Step 6.

Incubate the cells with the transfection complexes under their normal growth conditions, and monitor gene silencing after 48h. Change the medium as required.