

Jan 28,
2020

Bench top CUT&RUN with antibodies-online™ CUT&RUN Sets V.4

Forked from [Bench top CUT&Tag](#)

[antibodies online GmbH](#)¹

¹antibodies-online

1 *Works for me* [dx.doi.org/10.17504/protocols.io.bbteinje](https://doi.org/10.17504/protocols.io.bbteinje)

antibodies-online

Tech. support phone: **+49 0241 95163153** email: support@antibodies-online.com



Antibodies Online GmbH
antibodies online GmbH



ABSTRACT

This protocol is about Maintenance and Expansion of induced pluripotent stem cells.

EXTERNAL LINK

https://campaigns.antibodies-online.com/cutandrun-sets/?utm_source=protocolsio&utm_medium=protocol&utm_content=cutrun

THIS PROTOCOL ACCOMPANIES THE FOLLOWING PUBLICATION

The protocol is based on Skene; Henikoff JG; Henikoff S (2018): "[Targeted in situ genome-wide profiling with high efficiency for low cell numbers](#)"

GUIDELINES

Frequently Asked Questions

- [Why is the DNA yield so low?](#)
- [How can I validate that my primary antibody is working for CUT&RUN?](#)
- [Why do I need a negative control antibody? Why not just use a no-antibody control?](#)
- [Can I replace the antibody negative control using a knock-out \(or knock-down\) of my protein?](#)
- [Do I need to use a secondary antibody? Other CUT&RUN protocols do not use a secondary.](#)
- [Should I include heterologous spike-in DNA for quantitation?](#)
- [Are there protocols to make this work for tissue, where I cannot necessarily isolate single cells?](#)
- [Is it possible to use the CUT&RUN product sets with plant tissue samples?](#)
- [Can CUT&RUN be adapted for RIP-seq?](#)
- [Is it possible to use a cell free extract of a bacterial cell instead of immobilizing and permeabilizing the cell?](#)
- [Instead of the proteinase K digestion can I denature the proteins in the CUT&RUN product complexes by heat?](#)
- [What is preferable for DNA extractions prior to library preparation: phenol-chloroform extraction or affinity purification using a column?](#)

Why is the DNA yield so low?

CUT&RUN is performed using low cell numbers and the background signal is considerable lower than e.g. for ChIP. Due to these two factors the amount of recovered DNA is often times too low to be reliably measured based on a fluorometric assay or by capillary electrophoresis. PCR amplification of small CUT&RUN products, i.e. less than 50 bp, can be problematic and is therefore not any option. In order to assess the success of the CUT&RUN method each of the antibodies-online CUT&RUN product sets includes the CUT&RUN Positive Control antibody [ABIN6923144](#) against the abundant H3K27me3 histone modification. DNA fragments prepared using this antibody can be measured by sensitive electrophoresis on a Bioanalyzer or TapeStation or fluorometrically on a Qubit or Nanodrop fluorometer. When using the CUT&RUN Positive Control antibody [ABIN6923144](#) (or any other antibody specific for nucleosomal markers) a ladder corresponding to multiples of the 147 bp long nucleosomes should be visible by capillary electrophoresis.

How can I validate that my primary antibody is working for CUT&RUN?

For a CUT&RUN experiment the validation data could include e.g. a Tapestation or Bionalyzer plot showing the size distribution and qPCR data showing target enrichment.

As mentioned above, the DNA yield of an CUT&RUN experiment appears typically very low compared e.g. to ChIP-seq because of the lower initial sample size and the substantially lower DNA background. In particular for less abundant target protein the concentration is often times too low to be reliably measured using a fluorometric assay or by capillary electrophoresis. PCR amplification of small CUT&RUN products, i.e. less than 50 bp, can be problematic and is therefore not an option.

Once a sequencing library has been generated and sequenced map sequencing reads and verify the accumulation of reads at known binding sites.

Why do I need a negative control antibody? Why not just use a no-antibody control?

MNase is an endo- and exonuclease that will unspecifically bind and cleave unprotected DNA in hyper-accessible DNA, e.g. in regions surrounding regulatory elements. Free MNase will preferentially cut DNA within these hyper-accessible regions, thus potentially causing false positives and increased background signal in general.

In order to avoid this undesired effect of untethered MNase, the chromatin is randomly coated with the CUT&RUN Negative Control ([ABIN6923140](#)) prior to the addition of pA/G-MNase is added to the samples. pA/G-MNase is then tethered via its Protein A or Protein G portion to the antibodies' Fc fragment and background DNA fragmentation is dictated by the random antibody binding as opposed to the nuclease digestion of hyper-accessible DNA regions.

Can I replace the antibody negative control using a knock-out (or knock-down) of my protein?

Both controls are useful but address different aspects of the experiment and are therefore not interchangeable.

The CUT&RUN Negative Control ([ABIN6923140](#)) antibody is used to establish a reference background for peak calling. This is necessary because of the sparse background signal in CUT&RUN samples compared to ChIP-seq samples. The ko (or kd) control on the other hand gives an impression of unspecific binding of the antibody specific for the protein of interest to other proteins. It is useful to avoid identification of false positive signals.

Do I need to use a secondary antibody? Other CUT&RUN protocols do not use a secondary.

Depending on the host species and isotype of the antibody and the Protein A and/or Protein G MNase fusion protein, a secondary antibody may be necessary for pA/G-MNase binding.

Protein A has good high affinity to all rabbit IgG antibodies but low affinity to rat, goat and sheep IgG isotype antibodies and certain mouse IgG antibody subclasses, in particular IgG1. Protein G on the other hand binds well to the Fc region of mouse, goat, sheep, and most rat IgG. Its affinity to rabbit IgG however is lower than that of Protein A. When using pAG-MNase introduced with the improved CUT&RUN protocol it is therefore generally not necessary to use a secondary antibody. Use of the pA-MNase of the original protocol however might require the use of a secondary antibody raised in rabbit to assure efficient binding of the fusion protein to the antibody.

Should I include heterologous spike-in DNA for quantitation?

Our protocol is largely based on the improved CUT&RUN protocol. Here, the authors show that accurate quantitation is possible using heterologous spike-in DNA or carry-over *E. coli* DNA from the pA/G-MNase purification.

Are there protocols to make this work for tissue, where I cannot necessarily isolate single cells?

The Henikoff lab that published the original CUT&RUN method in 2017 has since developed several variations. One of these modification is AutoCUT&RUN which outlines a workflow to automate CUT&RUN for high-throughput characterization of single cells but also patient samples and xenografts.

In short, the tissue is processed manually or enzymatically into single, intact cell that are subsequently bound to ConA beads. The improved (and standard) CUT&RUN protocol that serve as basis for the protocol outlined in our CUT&RUN handbook can be adapted accordingly. Presently there is no work published documenting the use of intact tissue.

Is it possible to use the CUT&RUN product sets with plant tissue samples?

The CUT&RUN method can be applied without a problem to plant tissue samples (see e.g. [PMID30719569](#)). One essential step in addition to those lined out in the protocol is the generation of spheroblasts so that it becomes possible to permeabilize the plasma membrane for the application of the antibodies and the MNase fusion protein.

The positive control H3K27me3 antibody and the negative control guinea pig anti-rabbit IgG antibody as well as the ConA beads are suitable for use with plant samples. The antibodies that are included in some of the [sets](#), such as the anti-DYKDDDDK antibodies or the secondary antibodies, can also be used with plant samples. Whether they are needed or not depends on your experiment.

Can CUT&RUN be adapted for RIP-seq?

It should be possible to adapt the CUT&RUN protocol for use with RNA as an alternative to RIP-seq. RNA in the cytoplasm will attract the degradation machinery if it is lacking the 5' cap and the 3' poly-A tail. Thus, it is advisable to use isolated nuclei as sample material.

Consequently, no digitonin in the different buffers is needed since the nuclear envelope does not contain cholesterol. Isolated nuclei may be immobilized via glycoproteins on the nuclear envelope to the ConA beads like for a CUT&RUN experiment. An antibody against the protein of interest is added and subsequently the pA/G-MNase is tethered to the antibody, thus bringing the MNase into proximity of the RNA of interest. Similar to RIP-seq, isolated RNA is then translated into cDNA and can be sequenced and mapped.

Is it possible to use a cell free extract of a bacterial cell instead of immobilizing and permeabilizing the cell?

It should indeed be possible to use a bacterial extract as starting material instead of immobilized eukaryotic cells. There are a couple of things that should be considered:

One of the factors that makes CUT&RUN so much better than ChIP-seq in a eukaryotic system is that it is done in situ. The intact cells (or isolated nuclei) retain the large, undigested chromatin. Consequently, much fewer off-sites are present in the prepared DNA which is one of the reasons why CUT&RUN data sets have so much less background signal when compared to ChIP-seq data. This enrichment step will be missing when using a cell lysate as starting material. The CUT&RUN result should still be superior to the ChIP-seq result for the same starting material. However, it will be more "noisy" than CUT&RUN sequencing data from intact eukaryotic cells/nuclei.

The buffers described in the CUT&RUN protocol are designed to maintain the cells in the samples intact. The binding buffer contains some divalent cations that are necessary for the binding of the cells to the Concanavalin on the beads that are used for the immobilization. Subsequently, this buffer is changed for an antibody buffer that contains EDTA to chelate the divalent cations and prevent DNA cleavage prior to pA/G-MNase antibody binding. Since you are working with cell lysates as opposed to intact cells there is no need to include digitonin in any of the buffers. You don't need the binding buffer used to immobilize cells on the ConA beads. In order to control premature DNA cleavage by the MNase I would include some EDTA and/or EGTA in the lysis buffer.

Probably, you will have to titrate the amount of pA/G-MNase and try different digestion times to find the best balance between efficient cleavage of the intended sites and too much off-site cleavage.

The original CUT&RUN protocol foresees the use of heterologous spike-in DNA, e.g. from *E. coli*. In its latest iteration it is argued that there is no need to add this DNA since the pA/G-MNase preparation contains sufficient levels of *E. coli* DNA to serve as a standard for quantitation. In a prokaryotic system, spike-in DNA from a eukaryotic organism (e.g. *S. cerevisiae*) is recommended. Also keep in mind that the *E. coli* DNA carried over with the pA/G-MNase preparation may contain sequences that show homology to the bindings sites of your protein of interest.

Our product sets are optimized for use with eukaryotic cells. The positive control in all sets is a recombinant rabbit anti-H3K27me3 antibody. This control is not suitable for use with prokaryotic samples.

Instead of the proteinase K digestion can I denature the proteins in the CUT&RUN product complexes by heat?

Proteinase K digest followed by a phenol-chloroform extraction has historically been the go-to method to prepare high-molecular weight DNA, primarily from tissues. Subsequently to the diffusion of the CUT&RUN products out of the cells the DNA is present already in a relatively clean form, in particular when compared to a cell lysate like in ChIP-seq. Therefore, it is reasonable to assume that it is possible to forgo the Proteinase K treatment of the cells and instead proceed with the PCI extraction. As a reference, nucleic acid preparation using TRIzol does not include a Proteinase K digest. However, it has also been reported, that a proteinase K digest improves subsequent PCR amplification. Regarding heat inactivation instead of proteinase K digest in step, we recommend against this option. Your DNA of interest is at this point present in a complex consisting of the DNA, your antigen, the corresponding antibody, and the pA/G-MNase. If at this stage you boil your sample denature proteins you risk precipitation of your DNA with the protein compounds in this complex upon denaturation. This will also primarily affect the short CUT&RUN products, thus decreasing the signal to noise ratio in your library and potentially also reducing the library's complexity.

What is preferable for DNA extractions prior to library preparation: phenol-chloroform extraction or affinity purification using a column?

In the original publication describing the CUT&Tag method the authors mention the use of AMPure XP beads for the purification of the DNA subsequently to tagmentation and Proteinase K digest. A potential issue is the carry-over of active Proteinase K, which can interfere with the downstream PCR amplification. Therefore, the authors recommend now the phenol-chloroform extraction to assure complete denaturation of Proteinase K.

MATERIALS

NAME ▾	CATALOG # ▾	VENDOR ▾
antibodies-online™ CUT&RUN Pro Complete Set	ABIN6923135	antibodies-online
antibodies-online™ CUT&RUN Negative Control	ABIN6923140	antibodies-online
antibodies-online™ CUT&RUN Positive Control	ABIN6923144	antibodies-online

NAME ▾	CATALOG # ▾	VENDOR ▾
antibodies-online™ CUT&RUN Concanavalin A Beads	ABIN6923139	antibodies-online
antibodies-online™ CUT&RUN Secondary	ABIN6923141	antibodies-online
0.5 M EDTA pH 8.0	ABIN925554	antibodies-online
Trypan Blue (0.4%)	ABIN413910	antibodies-online
Proteinase K	ABIN920948	antibodies-online
CUT&RUN Pro Sec Set	ABIN6923137	antibodies-online
Spike-in Chromatin	ABIN4889666	antibodies-online
CUTANA™ pAG-MNase for ChIC/CUT&RUN Assays	ABIN6950951	antibodies-online

STEPS MATERIALS

NAME ▾	CATALOG # ▾	VENDOR ▾
antibodies-online™ CUT&RUN Concanavalin A Beads	ABIN6923139	antibodies-online
antibodies-online™ CUT&RUN Positive Control	ABIN6923144	antibodies-online
antibodies-online™ CUT&RUN Negative Control	ABIN6923140	antibodies-online
antibodies-online™ CUT&RUN Secondary	ABIN6923141	antibodies-online
CUTANA™ pAG-MNase for ChIC/CUT&RUN Assays	ABIN6950951	antibodies-online
antibodies-online™ CUT&RUN Concanavalin A Beads	ABIN6923139	antibodies-online
antibodies-online™ CUT&RUN Positive Control	ABIN6923144	antibodies-online
antibodies-online™ CUT&RUN Negative Control	ABIN6923140	antibodies-online
antibodies-online™ CUT&RUN Secondary	ABIN6923141	antibodies-online
antibodies-online™ CUT&RUN Secondary	ABIN6923141	antibodies-online
antibodies-online™ CUT&RUN Concanavalin A Beads	ABIN6923139	antibodies-online
antibodies-online™ CUT&RUN Positive Control	ABIN6923144	antibodies-online
antibodies-online™ CUT&RUN Negative Control	ABIN6923140	antibodies-online
antibodies-online™ CUT&RUN Secondary	ABIN6923141	antibodies-online
antibodies-online™ CUT&RUN Secondary	ABIN6923141	antibodies-online

MATERIALS TEXT

- DMEM/F12 medium
- Matrigel
- mTesR1 media
- Rock Inhibitor
- 1x PBS
- Accutase (Gibco A11105-01)



StemPro™ Accutase™ Cell Dissociation Reagent
by Thermo Fisher Scientific
Catalog #: A1110501

- 2X Freezing Medium (20 % DMXO, FBS)

SAFETY WARNINGS

Please see SDS (Safety Data Sheet) for hazards and safety warnings.

BEFORE STARTING



Warm all media before use. Media should be **Room temperature**. Do not use a water bath to warm media.

REAGENT SETUP (for 8 samples)

1

» Wash buffer (110 mL)

Component	Volume	Final concentration
ddH ₂ O	103 mL	-
1 M HEPES pH 7.5	2.2 mL	20 mM
5 M NaCl	3.3 mL	150 mM
2 M Spermidine	27.5 µL	0.5 mM

- Store Wash Buffer without protease inhibitors for up to one week at 4 °C.
- Add protease inhibitors fresh before use, e.g.:
1.1 mL Protease Inhibitor Cocktail 100x

» Binding Buffer (30 mL)

Component	Volume	Final concentration
ddH ₂ O	17.5 mL	-
1 M HEPES pH 7.5	360 µL	20 mM
2 M Spermidine	4,5 µL	0.5 mM
5% Digitonin	180 µL	0.05 %

- Store Antibody Buffer for up to one day at 4 °C until use.

» Digitonin Wash Buffer (55 mL)

Component	Volume	Final concentration
ddH ₂ O	17.5 mL	-
1 M HEPES pH 7.5	360 µL	20 mM
2 M Spermidine	4,5 µL	0.5 mM
5% Digitonin	180 µL	0.05 %

- Store Antibody Buffer for up to one day at 4 °C until use.

» Antibody Buffer (1 mL)

Component	Volume	Final concentration
ddH ₂ O	17.5 mL	-
1 M HEPES pH 7.5	360 µL	20 mM
2 M Spermidine	4,5 µL	0.5 mM
5% Digitonin	180 µL	0.05 %

- Store Antibody Buffer for up to one day at 4 °C until use.

» Low Salt Rinse Buffer (18 mL)

Component	Volume	Final concentration
ddH ₂ O	17.5 mL	-
1 M HEPES pH 7.5	360 µL	20 mM
2 M Spermidine	4,5 µL	0.5 mM
5% Digitonin	180 µL	0.05 %

• Store Antibody Buffer for up to one day at 4 °C until use.

» Low Salt Incubation Buffer (2 mL)

Component	Volume	Final concentration
ddH ₂ O	1953 µL	-
1 M HEPES pH 7.5	7 µL	3.5 mM
1 M CaCl ₂	20 µL	10 mM
5% Digitonin	20 µL	0.05%

• Store Low Salt Incubation Buffer for up to one week at 4 °C until use.

» Low Salt Stop Buffer (2 mL)

Component	Volume	Final concentration
ddH ₂ O	1700 mL	-
5 M NaCl	68 µL	170 mM
0.2 M EGTA	200 µL	20 mM








Store Low Salt Stop Buffer at 4 °C until use.

Add fresh before use:


Component	Volume	Final concentration
5% Digitonin	20 µL	0.05%
RNase A (10 mg/mL)	10 µL	50 µg/mL
Glycogen (20 mg/mL)	2.5 µL	25 µg/mL
Optional step: heterologous spike-in DNA		
heterologous spike-in DNA	-	100 pg/mL


I. Cell Harvest – at room temperature







- Harvest 10,000 to 500,000 cells for each sample at **Room temperature**. Keep cells for each sample in separate tubes.
- Centrifuge cell solution **00:03:00** at **600 x g** at **Room temperature**. Remove the liquid carefully.

- 4 Gently resuspend cells in  **1 ml Wash Buffer** by pipetting and transfer cell solution to a  **1.5 ml** microcentrifuge tube.
- 5 Centrifuge cell solution  **00:03:00** at  **600 x g** at  **Room temperature** and discard the supernatant.
- 6 Repeat steps 4-5 thrice for a total of four washes.  **go to step #4**
- 7 Resuspend cell pellet for each sample in  **1 ml Wash Buffer** by gently pipetting.

II. Concanavalin A beads preparation

- 8 Prepare one  **1.5 ml** microcentrifuge tube for each sample.
- 9 Gently resuspend the
- 10

 antibodies-online™ CUT&RUN
Concanavalin A Beads
by antibodies-online
Catalog #: ABIN6923139

Pipette  **10 µl CUT&RUN Concanavalin A Beads** slurry for each sample into the  **1.5 ml** microcentrifuge tubes.
- 11 Place the tubes on a magnet stand until the fluid is clear. Remove the liquid carefully.
- 12 Remove the microcentrifuge tube from the magnet stand.
- 13 Pipette  **1 ml Binding Buffer** into each tube and resuspend **CUT&RUN Concanavalin A Beads** by gentle pipetting.
- 14 Spin down the liquid from the lid with a quick pulse in a table-top centrifuge (max  **100 x g**).
- 15 Place the tubes on a magnet stand until the fluid is clear. Remove the liquid carefully.
- 16 Remove the microcentrifuge tube from the magnet stand.
- 17 Repeat steps 13-16  **go to step #13** twice for a total of three washes.
- 18 Gently resuspend the **CUT&RUN Concanavalin A Beads** in a volume of **Binding Buffer** corresponding to the original volume of bead slurry, i.e.  **10 µl** per sample.

III. Cell immobilization – binding to Concanavalin A beads

- 19 Carefully vortex the cell suspension from step [go to step #7](#) and add [10 µl](#) of the **CUT&RUN Concanavalin A Beads** in **Binding Buffer** prepared in **section II** to each sample.
- 20 Close tubes tightly and rotate for [00:05:00](#) to [00:10:00](#) at [Room temperature](#).

IV. Cell permeabilization and primary antibody binding

- 21 Place the microcentrifuge tubes on a magnet stand until the fluid is clear. Remove the liquid carefully.
- 22 Remove the microcentrifuge tubes from the magnet stand.
- 23 Place each tube at a low angle on the vortex mixer set to a low speed (approximately [1100 rpm](#)) and add [100 µl](#) **Antibody Buffer** containing **digitonin**.
- 24 Gently vortex the microcentrifuge tubes until the beads are resuspended.
- 25 Add [1 µl](#) **primary antibody** - against your antigen of interest, the



antibodies-online™ CUT&RUN Positive Control
by antibodies-online
Catalog #: ABIN6923144




, and the



antibodies-online™ CUT&RUN Negative Control
by antibodies-online
Catalog #: ABIN6923140

- corresponding to a 1:100 dilution (or a volume corresponding to the manufacturer's recommended dilution for immunofluorescence).

- 26 Rotate the microcentrifuge tubes for [00:05:00](#) to [00:10:00](#) at [Room temperature](#) or [02:00:00](#) to [Overnight](#) at [4 °C](#).
- 27 Spin down the liquid and place the tubes on a magnet stand until the fluid is clear. Remove the liquid carefully.
- 28 Remove the microcentrifuge tubes from the magnet stand.

- 29 Resuspend with  **1 ml Digitonin Wash Buffer** and mix by inversion. If clumping occurs, gently remove the clumps with a  **1 ml** pipette tip.
- 30 Repeat steps 27-29  **go to step #27** once for a total of two washes.



If no secondary antibody is used proceed directly to section VI. pA/G-MNase binding (Step 41).

V. Secondary antibody binding (optional)

31 



The



antibodies-online™ CUT&RUN



Secondary

by antibodies-online

Catalog #: ABIN6923141

is raised in rabbit against mouse IgG (H&L). In case a **primary antibody** from a different host species or isotype is used a different **secondary** must be selected accordingly.

Place the tubes on a magnet stand until the fluid is clear. Remove the liquid carefully.

- 32 Remove the microcentrifuge tubes from the magnet stand.
- 33 Vortex the sample at low speed (approximately  **1100 rpm**) and add  **100 µl Digitonin Wash Buffer** per sample along the side of the tube.
- 34 Tap to remove the remaining beads from the tube side.

35 Add  1 μ l




antibodies-online™ CUT&RUN
Secondary
by antibodies-online
Catalog #: ABIN6923141

corresponding to a 1:100 dilution (or a volume corresponding to the manufacturer's recommended dilution for immunofluorescence).

36 Rotate the microcentrifuge tubes for  01:00:00 at  4 °C .

37 Spin down the liquid and place the tubes on a magnet stand until the fluid is clear. Remove the liquid carefully.

38 Remove the microcentrifuge tubes from the magnet stand.




39 Resuspend with  1 ml **Digitonin Wash Buffer** and mix by inversion. If clumping occurs, gently remove the clumps with a 1 mL pipette tip.

40 Repeat steps 37-39  **go to step #33** once for a total of two washes.

VI. Protein A-MNase or Protein AG-MNase Binding

41 Place the tubes on a magnet stand until the fluid is clear. Remove the liquid carefully.




42 Remove the microcentrifuge tubes from the magnetic stand.



43 Vortex the sample at low speed (approximately  1100 rpm) and add  50 μ l **Digitonin Wash Buffer** per sample along the side of the tube. Add  2.5 μ l





CUTANA™ pAG-MNase for
ChIC/CUT&RUN Assays
by antibodies-online
Catalog #: ABIN6950951

Alternatively:













Vortex the sample at low speed (approximately  1100 rpm) and add  150 μ l **Digitonin Wash Buffer** containing  700 ng/mL of your own **pA/G-MNase** preparation per sample along the side of the tube.

44 Rotate the microcentrifuge tubes for  01:00:00 at  4 °C .

45 Spin down the liquid and place the tubes on a magnet stand until the fluid is clear. Remove the liquid carefully.

- 46 Remove the microcentrifuge tubes from the magnetic stand.
- 47 Resuspend with  **1 ml Digitonin Wash Buffer** and mix by inversion. If clumping occurs, gently remove the clumps with a 1 ml pipette tip.
- 48 Repeat steps 45-47  **go to step #45** once for a total of two washes.

VII. MNase digestion and release of pA/G bound chromatin fragments

- 49 Spin down the liquid from the lid with a quick pulse in a table-top centrifuge (max  **100 x g**).
- 50 Place the tubes on a magnet stand until the fluid is clear. Remove the liquid carefully.
- 51 Resuspend with  **1 ml Low Salt Rinse Buffer** and mix by inversion. If clumping occurs, gently remove the clumps with a 1 mL pipette tip.
- 52 Spin down the liquid from the lid with a quick pulse in a table-top centrifuge (max  **100 x g**).
- 53 Place the tubes on a magnet stand until the fluid is clear. Remove the liquid carefully.
- 54 Repeat steps 51-53  **go to step #51** once for a total of two washes.
- 55 Place each tube at a low angle on the vortex mixer set to a low speed (approx.  **1100 rpm**) and add  **200 µl ice cold Low Salt Incubation Buffer** per sample along the side of the tube.
- 56 Incubate tubes at  **0 °C** for the desired time (default is  **00:05:00**).
- 57 Place the tubes on a cold magnet stand until the fluid is clear. Remove the liquid carefully.
- 58 Remove the microcentrifuge tubes from the magnet stand.
- 59 Resuspend with  **200 µl Low Salt Stop Solution** and mix by gentle vortexing.
- 60 Incubate tubes at  **37 °C** for  **00:30:00** .
- 61 Place the tubes on a magnet stand until the fluid is clear.
- 62 Transfer the supernatant containing the pA/G-MNase-bound digested chromatin fragments to fresh  **1.5 ml** microcentrifuge tubes.


VIII. DNA extraction

- 63 Add **2 µl** 10% **SDS** to a final concentration of 0.1% and **5 µl** Proteinase K (**10 mg/mL**) to a final concentration of **2.5 mg/mL** to each supernatant from step 62.
- 64 Gently vortex tubes at a low speed of approximately **1100 rpm**.
- 65 Incubate tubes at **50 °C** for **01:00:00** or at **37 °C** **Overnight**
- 66 Add **200 µl** **PCI** to tube.
- 67 Vortex tubes thoroughly at high speed until the liquid appears milky.
- 68 Optional: Transfer liquid to a phase-lock tube.
- 69 Centrifuge tubes in a table-top centrifuge at **16000 x g** at **4 °C** for **00:05:00**.
- 70 Carefully transfer the upper aqueous phase to a fresh **1.5 ml** microcentrifuge tube containing **200 µl** **Chloroform:Isoamyl Alcohol 24:1**.
- 71 Vortex tubes thoroughly at high speed until the liquid appears milky.
- 72 Centrifuge tubes in a table-top centrifuge at **16000 x g** at **4 °C** for **00:05:00**.
- 73 Carefully transfer the upper aqueous phase to a fresh **1.5 ml** microcentrifuge tube containing **2 µl** glycogen (diluted 1:10 to **2 mg/mL** from the **20 mg/mL** stock solution).
- 74 Add **20 µl** **3 Molarity (M) NaOAc** or **100 µl** **5 Molarity (M) NH₄OAc**
- 75 Add **500 µl** **100% ethanol**.
- 76 Place tubes for **00:10:00** in a **dry ice/Ethanol mix** or **Overnight** at **-20 °C**.
- 77 Centrifuge tubes in a table-top centrifuge at **16000 x g** at **4 °C** for **00:05:00**.
- 78 Remove the liquid carefully with a pipette.
- 79 Add **1 ml** **70% ethanol**.

- 80 Centrifuge tubes in a table-top centrifuge at **16000 x g** at **4 °C** for **00:01:00**.
- 81 Remove the liquid carefully with a pipette.
- 82 Air-dry the pellet or dry the pellet in a SpeedVac.
- 83 Dissolve the pellet in **30 µl** **1 Milimolar (mM)** Tris-HCl, **0.1 Milimolar (mM)** EDTA.


IX. Sample quality control

- 84 Size distribution and concentration of the CUT&RUN products can be assessed at this point, e.g. using a



Qubit
Fluorometer

Invitrogen Q33228 [↗](#)



or



NanoDrop™ 3300 Fluorespectrometer
Fluorespectrometer

NanoDrop™ 3300 nd-3300 [↗](#)



or a Bioanalyzer or Tapestation. It is possible that the concentration of the recovered DNA is below the instrument's detection limit. It is also to be expected that the extracted DNA includes some large DNA fragments that will mask the signal of the CUT&RUN products. In this case it may be useful to PCR-amplify the DNA and check the library on a Bioanalyzer or Tapestation.

X. Sequencing library preparation

- 85 Prepare the CUT&RUN products sequencing libraries according to your established workflow. Because of the very low background with CUT&RUN, typically 5 million paired-end reads suffice for epitopes with a multitude of genomic binding sites, e.g. transcription factors or nucleosome modifications.

XI. Peak calling

- 86 The sparse background signal in CUT&RUN samples compared to ChIP-seq samples represents challenge for peak callers that employ statistical models relying on a high sequencing depth and high recall to identify true positives and avoid false positives. In contrast, peak calling for CUT&RUN data sets requires high specificity for true signal peaks. To this end, the Henikoff lab developed the Sparse Enrichments analysis for CUT&RUN (SEACR) peak caller that can be easily accessed using their web server at <https://seacr.fredhutch.org/>. Alternatively, the Orkin and Yuan labs have streamlined processing of CUT&RUN data using their CUT&RUNTools pipeline <https://bitbucket.org/qzhudfci/cutruntools/>



This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited