



Larval development 👄

PLOS Genetics

Wen Aw1

¹z3314717@unsw.edu.au

dx.doi.org/10.17504/protocols.io.rqzd5x6

Cage Studies



Wen Aw



EXT ERNAL LINK

https://doi.org/10.1371/journal.pgen.1007735

THIS PROTOCOL ACCOMPANIES THE FOLLOWING PUBLICATION

Aw WC, Towarnicki SG, Melvin RG, Youngson NA, Garvin MR, Hu Y, Nielsen S, Thomas T, Pickford R, Bustamante S, Vila-Sanjurjo A, Smyth GK, Ballard JWO (2018) Genotype to phenotype: Diet-by-mitochondrial DNA haplotype interactions drive metabolic flexibility and organismal fitness. PLoS Genet 14(11): e1007735. doi: 10.1371/journal.pgen.1007735

PROTOCOL STATUS

Working

- Flies from each of the 4 mitotypes were placed in individual egg collection containers.
- Eggs were collected manually and added to either the 1:2 or 1:16 P:C diet with ~200 eggs per bottle with 6 bottles/mitotype/diet.
- Microbiome was added after 48 hours and flies that eclosed in a 3d window were collected, counted and % eclosion was 3 determined by dividing the number of eclosed flies from each mitotype by the sum of both mitotypes.

This is an open access protocol distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited