



Mar 28,
2019

Working

E gene amplification

Bo Yi¹, Yi Chen¹, Xiao Ma¹, Haibin Wang², Rong Wang¹, Keqin Ding¹, Lei Xie¹, Dongliang Zhang¹, Shuli Jiao¹, Xuying Lao¹, Yi-Chen Chiang³, Yanhua Su³, Benhua Zhao³, Guozhang Xu¹, Tianmu Chen³

¹Ningbo Municipal Center for Disease Control and Prevention, ²Haishu District Center for Disease Control and Prevention, ³State Key Laboratory of Molecular Vaccinology and Molecular Diagnostics, School of Public Health, Xiamen University

[dx.doi.org/10.17504/protocols.io.zkjf4un](https://doi.org/10.17504/protocols.io.zkjf4un)

 Tianmu Chen 

PROTOCOL STATUS

Working

We use this protocol in our group and it is working

- 1 The dengue virus strains obtained from cell culture were extracted by TGuide S32 magnetic bead method DNA/RNA extraction kit of Tiangen Biochemical Technology (Beijing) Co., Ltd. and amplified by one step RNA PCR Kit (Code No: DRR057A) reagent of Bao Bioengineering (Dalian) Co., Ltd.
- 2 The full sequence of E gene amplified by primer sequence was D1 (E) F: CAA GAA CCG AAA CA / GT GGA TGT C; D1 (E) R: GGC TGA TCG AAT TCC ACA CAC, and the length of amplified fragment was 1849 bp.
- 3 Reaction conditions: reverse transcription at 50 °C 30 min, reverse transcription at 94°C 2 min, reverse transcription at 94°C 30 s, reverse transcription at 52°C 30 s, reverse transcription at 72°C 2 min, 40 cycles and extension at 72°C. The amplified product was 5 micron L, and 1.5% agarose gel electrophoresis was used to confirm the reaction product according to Marker position.



This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited