



Feb 06, 2020

Enzymatic fingerprinting of AX and mixed-linked BG [↗](#)

PLOS One

Alison Lovegrove<sup>1,2</sup><sup>1, 2</sup>Rothamsted Research**1** Works for me [dx.doi.org/10.17504/protocols.io.babriam6](https://doi.org/10.17504/protocols.io.babriam6)

Alison Lovegrove

## ABSTRACT

Lab protocol for: Enzymatic fingerprinting for arabinoxylan and mixed-linked beta-glucan in wheat flour by HPAEC-PAD. Based on Ordaz-Ortiz and Saulnier, 2005 J. Cereal Sci. 42. 119-125. and Odaz-Ortiz et al., 2005 J. Ag Food Chem. 53. 8349-8356.

## EXTERNAL LINK

<https://doi.org/10.1371/journal.pone.0227826>

## THIS PROTOCOL ACCOMPANIES THE FOLLOWING PUBLICATION

Lovegrove A, Wingen LU, Plummer A, Wood A, Passmore D, Kosik O, Freeman J, Mitchell RAC, Hassall K, Ulker M, Tremmel-Bede K, Rakszegi M, Bedő Z, Perretant M, Charmet G, Pont C, Salse J, Waite ML, Orford S, Burrridge A, Pellny TK, Shewry PR, Griffiths S (2020) Identification of a major QTL and associated molecular marker for high arabinoxylan fibre in white wheat flour. PLoS ONE 15(2): e0227826. doi: [10.1371/journal.pone.0227826](https://doi.org/10.1371/journal.pone.0227826)

## ATTACHMENTS

[SOP. Fingerprinting of AX and MLG in wheat flour.docx](#)

This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited