



Jun 26,
2019

GG1 - sgRNA cloning for *Phaeodactylum tricornutum*

Mark Moosburner¹, Andrew Allen¹

¹University of California, San Diego

Working

dx.doi.org/10.17504/protocols.io.4abgsan

JCVI West Protocols



Mark Moosburner

Scripps Institution of Oceanography, J. Craig Venter Institu...



ABSTRACT

The CRISPR-Cas9 gene mutagenesis system was adapted for the marine diatom *Phaeodactylum tricornutum* (CCAP-1055/1) (Figure 1). Here, Cas9 and sgRNA(s) were delivered to *Phaeodactylum* by bacterial-conjugation transformation on an episome, or artificial chromosome, that is stably maintained and replicated independently from and with the *Phaeodactylum* chromosomal DNA. Two sgRNA expression cassettes, sgRNA (1/1) and sgRNA (1/2) were cloned into the episome. The sgRNAs each contain a unique 20-nucleotide spacer sequence that, after sgRNA cassette transcription, guides the Cas9 to a nucleic acid target by complementary binding followed by Cas9 nuclease activity. Two *Phaeodactylum* genes, Pt_GSII (Gene ID: 51092) and Pt_cGOGAT (Gene ID: 24739), were targeted by each sgRNA, respectively. The following protocol was followed to synthesize spacer sequences and sgRNA expression cassettes



GG1_ Spacer Cloning for
P.t. episome.pdf



This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited