




Making OP50 solution from Frozen Stock

Priota Islam¹

¹Imperial College London

dx.doi.org/10.17504/protocols.io.vube6sn

Behavioural Genomics



PROTOCOL STATUS

Working

We use this protocol in our group and it is working

¹ Day: 1

- Get LB Agar from the media kitchen and autoclave it for 2hrs
- Post autoclave, pour the agar on large petri dishes (60mm) and leave to dry overnight to be transferred to the cold room the next day

² Day: 2

- Get the frozen tube out of the freezer (Freezer 12, SD Box)
- Take a sterile pipette tip and get some frozen sample
- Streak the pipette on the LB agar plate
- Take another new pipette and streak across the old streak and then at a separate spot streak again
- Repeat this step for one more time
- Keep the plate at 37°C incubator overnight for the bacteria to grow
- Transfer the plate to the fridge the next day (This plate can be used for about 1 month to inoculate bacterial culture)

³ Day:3

- Purchase LB Broth from the Media kitchen (Don't use less than 200ml)
- Get flat bottomed conical flasks from the glassware kitchen (For 200ml LB Broth take 400/500ml flask to allow enough headspace for the bacteria)
- From the culture plate, select and circle a single colony
- Using aseptic technique, carefully scoop the single colony and mix it with the LB Broth that is already poured into the conical flask
- Label the flask and put it on the 37°C shaker for overnight incubation

4 Day:4

- Following overnight incubation, take the flask out of the incubator and measure the OD using a spectrophotometer
- Record the average of the OD600, use LB Broth as Blank
- Aliquot the solution into labelled 15ml falcon tubes and keep them in the 4°C Fridge

This is an open access protocol distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited