



Jun 20,  
2019

## Fabrication of DNA constructs by Gibson Assembly and Golden Gate reactions V.2

Tamara Matute<sup>1</sup>, Isaac Nuñez<sup>1</sup>, Peter Von Dassow<sup>1</sup>, Fernan Federici<sup>1</sup>

<sup>1</sup>PUC

Working

dx.doi.org/10.17504/protocols.io.4fcgtiw

Protist Research to Optimize Tools in Genetics (PROT-G)

SynBioUC Chile



Fernan Federici



### ABSTRACT

We used this protocol to build a series of codon optimized BFP expressing vectors in the context of the EMS initiative from Gordon and Betty Moore foundation. A complete list of files corresponding to the DNA parts and vectors used in this study can be found at our [Drive](#) and [zenodo](#) database.

### Introduction and rationale

- 1 Golden Gate assembly creates vectors by combining vectors containing level 0 parts (e.g. promoters, CDSes, terminators) and an acceptor vector. The first step involves the creation of libraries of level 0 parts by 'domesticating' DNA sequences of interest (e.g a promoter or a new fluorescent protein).

### Domestication of DNA sequences into level 0 part vectors

- 2 Level 0 parts for Golden Gate reactions were generated by Gibson Assembly. Parts can be generated by PCRing natural sequences (e.g genomic DNA) or existing vectors; or by fully synthesising (e.g. gblocks from IDT) DNA fragments of interest.

For PCRed parts, reactions were performed with Phusion® High-Fidelity DNA Polymerase following this conditions:

#### Thermocycling

98°C x 30 sec

(98°C x 10 sec + 60°C x 30 sec + 72°C x 2 min) x 35 cycles

72°C x 10 min

4°C

#### PCR mix:

1ul primer forward Gibson

1ul primer reverse Gibson

1,33ul template (concentration: 1ng/ul)

8 ul Buffer HF 5X

0,8ul dNTPs (10uM)

0,4ul Phusion® High-Fidelity DNA Polymerase (2U/ul)

27,46 ul H2O

Commerically synthesised DNA fragments were ordered containing the 20-30bp overhangs used for Gibson assembly into the level 0 part vector.

### Gibson assembly step for the creation of level 0 part vector

- 3 Desired PCR products were identified by gel electrophoresis and extracted with the purification kit Wizard SV Gel & PCR Clean-Up System (Promega). DNA concentration were corroborated with a Take3™ Micro-Volume Synergy™ HTX before performing Gibson reactions. The volumes used for each part were calculated by eq 1, 2 and 3.

Eq.1

$$X [pMol/\mu l] = \frac{Concentration [ng/\mu l] * 1000}{650 [\frac{gr}{mol}] * Lenght [bp]}$$

Eq.2

$$Vol_{ratio\ i} = \frac{X}{pmol_{req}}$$

Volume ratio [ul] = pmol/required pmol

where required pmol is:

0,01 for parts >2500 bp

0,03 for parts >200 bp and <2500 bp

0,05 for parts <200 bp

Volume to be added was calculated as:

Eq.3

$$Vol_{piece\ i} = \frac{Vol_{total} * Vol_{ratio\ i}}{\sum_i^n Vol_{ratio\ i}}$$

1,5µl of parts were combined with 4,5µl of Gibson Master Mix in ice. This reaction was incubated at 50°C for 1 hour.

#### Reagents:

##### Gibson Mix (1.33X):

100 ul 5X Isothermal Buffer

2 ul T5 Exonuclease (1 U/ul)

6,25 ul Phusion® High-Fidelity DNA Polymerase (2U/ul)

50 ul Taq DNA ligase (40U/ul)

216,75 ul H2O

Store at -20°C.

**5X Isothermal Buffer:**

25% PEG-8000  
500 mM Tris-HCl pH 7,5  
50 mM MgCl<sub>2</sub>  
50 mM DTT  
1 mM dATP  
1 mM dTTP  
1 mM dCTP  
1 mM dGTP  
5 mM NAD  
H<sub>2</sub>O  
Store at -80°C

Transformation of Level 0 donor vectors

- 4 These reactions were transformed into chemically competent TOP10 cells prepared by the OOW protocol ([http://www.openwetware.org/wiki/TOP10\\_chemically\\_competent\\_cells](http://www.openwetware.org/wiki/TOP10_chemically_competent_cells)), miniprep and sequenced. These level 0 parts were stored as vectors for the following Golden Gate reactions.

Golden Gate assembly of promoters and fluorescent proteins combinations

- 5 Final combinatorial assemblies of promoters, CDSes and terminators here described were produced by Golden Gate. For this, a general acceptor vector was created that accepts level 0 parts for the assembly of transcriptional units (e.g. promoter-CDS-terminator of interest).

Eq.4

$$Vol_{required} [ul] = \frac{X * Length * 650}{[concentration] * 10^6}$$

The assembly was performed as follows:  
(37°C x 3 min + 16°C x 5 min) \* 40 cycles.  
50°C x 5 min.  
80°C x 10 min  
4°C

Transformation of Golden Gate reactions

- 6 3µl of these reactions were used to transform TOP10 Escherichia coli cells following the step described above.

Positive colonies were identified by colony PCR and sequenced.



This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited