



Sep 13, 2019

## MojoSort™ Isolation Kits Column Protocol - 3

Sam Li<sup>1</sup><sup>1</sup>BioLegend

1 Works for me

[dx.doi.org/10.17504/protocols.io.7behije](https://doi.org/10.17504/protocols.io.7behije)

BioLegend

Sam Li  
BioLegend

## ABSTRACT

BioLegend MojoSort™ nanobeads work in commonly used separation columns, based on our internal research as well as validation by external testing by academic labs. This simple protocol consists of following the MojoSort™ protocol to label the cells with **pre-diluted** MojoSort™ reagents and using the columns as indicated by the manufacturer.

**Note:** Due to the properties of our beads, it may be possible to use far fewer beads and less antibody cocktail than with other commercial suppliers. We recommend a titration to find the best dilution factor. However, as a general rule, dilutions ranging from 1:2 to 1:10 for the antibody cocktail can be used. Dilutions ranging from 1:5 to 1:20 for the Streptavidin Nanobeads can be used. Please contact BioLegend Technical Service ([tech@biolegend.com](mailto:tech@biolegend.com)) if further assistance is needed.

## EXTERNAL LINK

<https://www.biolegend.com/protocols/mojosort-isolation-kits-column-protocol-3/4591/>

## GUIDELINES

MojoSort™ magnetic particles can be used with other commercially available magnetic separators, both free standing magnets and column-based systems. Because MojoSort™ protocols are optimized for the MojoSort™ separator, the protocols may need to be adjusted for other systems. Please contact BioLegend Technical Service ([tech@biolegend.com](mailto:tech@biolegend.com)) for more information and guidance. We do not recommend using MojoSort™ particles for BD's IMag™ or Life Technologies' DynaMag™.

## MATERIALS

NAME ▾	CATALOG # ▾	VENDOR ▾
MojoSort™ Buffer	480017	BioLegend

## MATERIALS TEXT

Additional reagents:

- commercially available cell separation columns
- 5 mL polypropylene tubes
- 70 µm cell strainer

This protocol works with the following MojoSort™ Kits (cat#):

480003, 480004, 480011, 480012, 480129, 480065, 480066

- 1 Prepare cells from your tissue of interest or blood without lysing erythrocytes. Kits for human samples have been optimized for PBMCs, please prepare the cells using a suitable method.
- 2 In the final wash of your sample preparation, resuspend the cells in MojoSort™ Buffer by adding up to 4 mL in a 5 mL (12 x 75 mm) polypropylene tube.  
**Note:** Keep MojoSort™ Buffer on ice throughout the procedure.

- 3 Filter the cells with a 70  $\mu\text{m}$  cell strainer, centrifuge at 300xg for 5 minutes, and resuspend in an appropriate volume of MojoSort™ Buffer. Count and adjust the cell concentration to  $1 \times 10^8$  cells/mL. 5m
- 4 Aliquot 100  $\mu\text{L}$  ( $10^7$  cells) into a new tube. **Add 10  $\mu\text{L}$  of the pre-diluted Biotin-Antibody Cocktail**. Mix well and **incubate on ice for 15 minutes**. Scale up the volume if separating more cells. For example, add 100  $\mu\text{L}$  of pre-diluted antibody cocktail for separating  $1 \times 10^8$  cells in 1 ml of MojoSort™ Buffer. When working with less than  $10^7$  cells, use indicated volumes for  $10^7$  cells. 15m
- 5 Vortex the Streptavidin conjugated Nanobeads (to resuspend) at max speed, 5 touches, and prepare the dilutions to test. **Add 10  $\mu\text{L}$  of pre-diluted Streptavidin Nanobeads**. Mix well and **incubate on ice for 15 minutes**. Scale up the volume accordingly if separating more cells. For example, add 100  $\mu\text{L}$  of pre-diluted Nanobeads for separating  $1 \times 10^8$  cells in 1 ml of MojoSort™ Buffer. When working with less than  $10^7$  cells, use indicated volumes for  $10^7$  cells. 15m
- 6 Wash the cells by adding MojoSort™ Buffer up to 4 mL. Centrifuge the cells at 300xg for 5 minutes. 5m
- 7 Discard the supernatant.
- 8 Resuspend the cells in appropriate amount of MojoSort™ Buffer and proceed to separation. At least 500  $\mu\text{L}$  is needed for column separation.

**Note:** There are several types of commercially available columns, depending on your application. Choose the one that fits best your experiment:

	Max. number of labeled cells	Max. number of total cells	Cell suspension volume	Column rinse volume	Cell wash volume	Elution volume
<b>Small Capacity</b>	$1 \times 10^7$	$2 \times 10^8$	500 $\mu\text{L}$ for up to $10^8$ cells	1ml	1 ml	1 ml
<b>Medium Capacity</b>	$1 \times 10^8$	$2 \times 10^9$	500 $\mu\text{L}$ for up to $10^9$ cells	3ml	3 ml	5 ml
<b>Large Capacity</b>	$1 \times 10^9$	$2 \times 10^{10}$	500 $\mu\text{L}$ for up to $10^{10}$ cells	20-50ml	30 ml	20 ml

Example of magnetic separation with medium capacity columns:

- 9 Place the column in a magnetic separator that fits the column.
- 10 Rinse the column with 3 mL of cell separation buffer.
- 11 Add the labeled cell suspension in at least 500  $\mu\text{L}$  of buffer to the column through a 30  $\mu\text{m}$  filter and collect the fraction containing the unlabeled cells. These are the cells of interest; do not discard.
- 12 Wash the cells in the column **2 times** with 3 mL of buffer and collect the fraction containing the untouched cells. Combine with the collected fraction from step 3.

- 13 If desired, the labeled cells can be collected by taking away the column from the magnet and place it on a tube. Then add 5 mL of buffer and flush out the magnetically labeled fraction with a plunger or supplied device. The labeled cells may be useful as staining controls, to monitor purity/yield, or other purposes.



This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited