



Oct 06, 2019

## Phylogenetic analysis

Norfitriah Mohamed Sohaimi<sup>1</sup>, Mohd Hair Bejo<sup>1</sup>, Abdul Rahman Omar<sup>1</sup>, Aini Ideris<sup>1</sup>, Nurulfiza Mat Isa<sup>2</sup>

<sup>1</sup>Faculty of Veterinary Medicine, Universiti Putra Malaysia, Serdang, Selangor, Malaysia., <sup>2</sup>Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia, Serdang, Selangor, Malaysia.

1 Works for me dx.doi.org/10.17504/protocols.io.7zihp4e



### MATERIALS TEXT

Twenty-eight published FAdVs hexon (Table 1) and fiber (Table 2) gene sequences were obtained from GenBank. Another two reference strains from different genus avian adenovirus was used which are Duck adenovirus (Atadenovirus) and Turkey adenovirus (Siadenovirus). Multiple sequence alignment of nucleotide and deduced amino acid were conducted using BioEdit Version 7.2.5 and Mega Version 5 software. A region of L1 loop in hexon gene with 198 amino acids and entire region of fiber gene was selected for construction of phylogenetic tree. Jones-Taylor-Thorton (JTT) model was used to compute distance matrix using MEGA software and followed by phylogenetic tree construction using Neighbour-joining method with 1000 bootstrap replicates [Juliana et al., 2104].

☐ Table 1.docx ☐ Table 2.docx

1



This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited