



Oct 06, 2019

Propagation and attenuation of FAdV isolate in CEL cells

Norfitriah Mohamed Sohaimi¹, Mohd Hair Bejo¹, Abdul Rahman Omar¹, Aini Ideris¹, Nurulfiza Mat Isa²

¹Faculty of Veterinary Medicine, Universiti Putra Malaysia, Serdang, Selangor, Malaysia., ²Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia, Serdang, Selangor, Malaysia.

1 Works for me dx.doi.org/10.17504/protocols.io.7zdhp26



Norfitriah Mohamed Sohaimi

MATERIALS

NAME

CATALOG #

VENDOR


Fetal Bovine Serum

10270106

Gibco - Thermo Fischer

MATERIALS TEXT

A confluent monolayer was washed twice with serum free medium and inoculated with 0.1 mL of homogenate liver embryos and labeled as first passage (CEL1). Infected flasks were incubated at 37°C for 60 minutes for virus adsorption and added with maintenance medium containing 2% fetal bovine serum (FBS) under 37°C incubator. The cells were observed daily under inverted microscope for cytopathic effect (CPE) for 3 days post-inoculation (pi). Flasks with prominent CPE were harvested by 3 times repeated frozen and thawed prior centrifugation at 216 x *g* for 10 minutes. Virus supernatant was collected and stored at -20°C prior inoculation for subsequence passages. Fresh confluent monolayer was prepared for each passage and inoculated with 0.1 mL of viral supernatant and continued until 35th consecutive passage. For non-infected flasks, monolayer remained uninoculated and was used as control cells.



This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited