

Spot-on-lawn (halo) assay for screening enrichment cultures and isolates for viruses

Kenneth M. Stedman, Kate Porter, and Mike L. Dyall-Smith

Abstract

This protocol is based on Schleper et al. (1992) as modified by Stedman et al. (2003).

This is a protocol from:

Stedman, K. M., K. Porter, and M. L. Dyall-Smith. 2010. Chapter 6: The isolation of viruses infecting Archaea. Manual of Aquatic Viral Ecology. Waco, TX: American Society of Limnology and Oceanography. doi:10.4319/mave.2010.978-0-9845591-0-7

Please see the [published manuscript](#) for additional information.

Citation: Kenneth M. Stedman, Kate Porter, and Mike L. Dyall-Smith Spot-on-lawn (halo) assay for screening enrichment cultures and isolates for viruses. **protocols.io**

dx.doi.org/10.17504/protocols.io.eagbabw

Published: 09 Dec 2015

Guidelines

Authors: Kenneth M. Stedman¹, Kate Porter², and Mike L. Dyall-Smith³

¹Department of Biology, Center for Life in Extreme Environments, Portland State University, P.O. Box 751, Portland, OR 97207, USA

²Biota Holdings Limited, 10/585 Blackburn Road, Notting Hill Victoria 3168, Australia

³Max Planck Institute of Biochemistry, Department of Membrane Biochemistry, Am Klopferspitz 18, 82152 Martinsried, Germany




Fig. 1: Pictorial overview of isolation of *Sulfolobus* viruses.

(A) Wolfram Zillig sampling at a typical *Sulfolobus*-containing pool in Yellowstone National Park, USA, September 2000 (inset shows anaerobic tubes with samples).

(B) 80°C incubator with long-necked growth flasks (detail in inset).

(C) Single-colony isolates of *Sulfolobus solfataricus* on a Gelrite® plate. This plate contains a mixture of *S. solfataricus* containing (blue colonies) and lacking (brown) a vector expressing the *lacS* gene from *S. solfataricus* and was sprayed with 5-bromo-4-chloro-3-indolyl-b-D-galactopyranoside (X-gal) (see Jonuscheit et al. 2003).

(D) Lawn of *S.solfataricus* strain P1 with halos of growth inhibition due to virus production by 2- μ l spots of virus-infected strains. Spots labeled SV2P1 and SV2P2 are from *S. solfataricus* strains P1 and P2 infected with SSV-I2 respectively (Stedman et al. 2003). Spot labeled C is a detergent-positive control. Spot labeled P2- is uninfected *S. solfataricus* strain P2 as a negative control.

Protocol

Step 1.

Gelrite plates are preincubated ca. 10 min at 80°C to dry.

 DURATION

00:10:00

Step 2.

10 mL of *Sulfolobus* medium with ca. 0.2 % (w/v) Gelrite is boiled to dissolve the Gelrite.

Step 3.

This “softlayer” is allowed to cool slightly (to ca. 80°C).

Step 4.

Approximately 3 mL of softlayer are added to ca. 0.2 mL of exponentially growing host cells, generally *Sulfolobus solfataricus*, and spread on a plate by swirling.

Step 5.

After the Gelrite solidifies, 1–2 μ L of culture or supernatant to be screened is spotted on the plate.

Step 6.

For a positive control, 1 μ L of a 0.01% (v/v) Triton X-100 solution is spotted.

Step 7.

Plates are incubated as above for 2–3 d and plates examined for clearing around spots (Fig. 1D in guidelines).

 DURATION

12:00:00