



## Tranfection of SpCas9 containing plasmids to generate cell lines with Cas9

Binnypreet Kaur<sup>1, 2</sup>, Drahomíra Faktorová<sup>1, 2</sup>, Julius Lukeš<sup>1, 2, 3, 4</sup>

<sup>1</sup>Institute of Parasitology, Czech Academy of Sciences, <sup>2</sup>Faculty of Sciences, University of South Bohemia, České Budějovice, Czech Republic, <sup>3</sup>Institute of Parasitology, Czech Academy of Sciences, Czech Republic, <sup>4</sup>Canadian Institute for Advanced Research, Toronto, Canada

[dx.doi.org/10.17504/protocols.io.x36fqre](https://doi.org/10.17504/protocols.io.x36fqre)

 Binnypreet Kaur   

### ABSTRACT

We electropoarted Spcas9 containing plasmid (pVY015) with *Leishmania mexicana* UTR under hygromycin selection marker as described in Protocol 7.

### PROTOCOL STATUS

#### Working

We use this protocol in our group and it is working

- 1 We electropoarted Spcas9 containing plasmid (pVY015) with *Leishmania mexicana* UTR under hygromycinselection marker as described in Protocol 7.



This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited