

Restriction Digest Version 2

New England Biolabs

Abstract

The following is a "typical" restriction endonuclease reaction. Please see the "guidelines" tab below for the NEB tips on optimizing restriction digests.

Citation: New England Biolabs Restriction Digest. [protocols.io](https://www.protocols.io)

[dx.doi.org/10.17504/protocols.io.isycefww](https://doi.org/10.17504/protocols.io.isycefww)

Published: 02 Mar 2018

Guidelines

Guidelines for Optimizing Restriction Endonuclease Reactions

If you are using a Master Mix, see '[optimizing RE-Mix® reactions](#)'.

There are several key factors to consider when setting up a restriction endonuclease digest. Using the proper amounts of DNA, enzyme and buffer components in the correct reaction volume will allow you to achieve optimal digestion. By definition, 1 unit of restriction enzyme will completely digest 1 µg of substrate DNA in a 50 µl reaction in 60 minutes. This enzyme : DNA : reaction volume ratio can be used as a guide when designing reactions. However, most researchers follow the 'typical' reaction conditions listed, where a 5–10 fold overdigestion is recommended to overcome variability in DNA source, quantity and purity. NEB offers the following tips to help you to achieve maximal success in your restriction endonuclease reactions.

A 'Typical' Restriction Digest

Restriction Enzyme	10 units is sufficient, generally 1µl is used
DNA	1 µg
10X NEBuffer	5 µl (1X)
Total Reaction Volume	50 µl
Incubation Time	1 hour*
Incubation Temperature	Enzyme dependent

* Can be decreased to 5-15 minutes by using a [Time-Saver™ Qualified enzyme](#).

Enzyme

- Keep on ice when not in the freezer
- Should be the last component added to reaction
- Mix components by pipetting the reaction mixture up and down, or by 'flicking' the reaction tube.
- Follow with a quick ('touch') spin-down in a microcentrifuge.

- Do not vortex the reaction.
- In general, we recommend 5–10 units of enzyme per µg DNA, and 10–20 units for genomic DNA in a 1 hour digest.
- NEB has introduced a line of [High-Fidelity \(HF®\) enzymes](#) that provide added flexibility to reaction setup.

DNA

- Should be free of contaminants such as phenol, chloroform, alcohol, EDTA, detergents or excessive salts. Extra wash steps during purification are recommended.
- Methylation of DNA can inhibit digestion with certain enzymes. For more information about methylation, [Effect of CpG Methylation on Restriction Enzyme Cleavage](#) and [Dam and Dcm Methylases of *E.coli*](#)

Buffer

- Use at a 1X concentration
- Supplement with SAM (S-Adenosyl methionine) to the recommended concentration if required.

Reaction Volume

- A 50 µl reaction volume is recommended for digestion of 1 µg of substrate
- Enzyme volume should not exceed 10% of the total reaction volume to prevent [star activity](#) due to excess glycerol
- Additives in the restriction enzyme storage buffer (e.g., glycerol, salt) as well as contaminants found in the substrate solution (e.g., salt, EDTA, or alcohol) can be problematic in smaller reaction volumes. The following guidelines can be used for techniques that require smaller reaction volumes.

	Restriction Enzyme*	DNA	10X NEBuffer
10 µl rxn**	1 unit	0.1 µg	1 µl
25 µl rxn	5 units	0.5 µg	2.5 µl
50 µl rxn	10 units	1 µg	5 µl

* Restriction Enzymes can be diluted using the recommended diluent buffer when smaller amounts are needed.

** 10 µl rxns should not be incubated for longer than 1 hour to avoid evaporation.

Incubation Time

- Incubation time is typically 1 hour
- Can often be decreased by using an excess of enzyme, or by using one of our [Time-Saver Qualified enzymes](#).
- It is possible, with many enzymes, to use fewer units and digest for up to 16 hours. For more information, visit [Extended Digests with Restriction Endonucleases](#).

Stopping a Reaction

If no further manipulation of DNA is required:

- Terminate with a stop solution (10 µl per 50 µl rxn) [1x: 2.5% Ficoll®-400, 10mM EDTA, 3.3mM Tris-HCl, 0.08% SDS, 0.02% Dye 1, 0.001% Dye 2, pH 8.0@25°C] (e.g., NEB [#B7024](#))

When further manipulation of DNA is required:

- [Heat inactivation](#) can be used
- Remove enzyme by using a spin column or phenol/chloroform extraction

Storage

- Storage at -20°C is recommended for most restriction enzymes. For a few enzymes, storage at -70°C is recommended for periods longer than 30 days. Please refer to the enzyme's technical data sheet or catalog entry for storage information.
- 10X NEBuffers should also be stored at -20°C

Stability

All enzymes are assayed for activity every 4 months. The expiration date is found on the label. Exposure to temperatures above -20°C should be minimized whenever possible

Control Reactions

If you are having difficulty cleaving your DNA substrate, we recommend the following control reactions:

- Control DNA (DNA with multiple known sites for the enzyme, e.g. lambda or adenovirus-2 DNA) with restriction enzyme to test enzyme viability
- If the control DNA is cleaved and the experimental DNA resists cleavage, the two DNAs can be mixed to determine if an inhibitor is present in the experimental sample. If an inhibitor (often salt, EDTA or phenol) is present, the control DNA will not cut after mixing.

Protocol

Step 1.

Set up the following reaction (total reaction volume **50µl**).

Restriction Enzyme	10 units is sufficient, generally 1µl is used
DNA	1 µg
10X NEBuffer	5 µl (1X)
Total Reaction Volume	50 µl
Incubation Time	1 hour*
Incubation Temperature	Enzyme dependent

* Can be decreased to 5-15 minutes by using a [Time-Saver™ Qualified enzyme](#).

PROTOCOL

. [Restriction Digest Reaction](#)

CONTACT: [New England Biolabs](#)

NOTES

New England Biolabs 06 Oct 2014

Enzyme volume should not exceed 10% of the total reaction volume to prevent [star activity](#) due to excess glycerol.

New England Biolabs 06 Oct 2014

A 50 µl reaction volume is recommended for digestion of 1 µg of substrate.

New England Biolabs 08 Oct 2014

The enzyme should be the last component added to reaction

New England Biolabs 05 Jul 2017

Keep Enzyme on ice when not in freezer.

Step 1.1.

DNA **1µg**

AMOUNT

1 µg Additional info:

Step 1.2.

10X NEBuffer **5µl** (1X)

AMOUNT

5 µl Additional info:

Step 1.3.

Restriction Enzyme, 10 units is sufficient, generally **1µl** is used

Step 2.

Mix components by pipetting the reaction mixture up and down, or by "flicking" the reaction tube.

Step 3.

Quick ("touch") spin-down in a microcentrifuge. **Do not vortex the reaction.**

Step 4.

Incubate for 1 hour at the enzyme-specific appropriate temperature.

📌 **NOTES**

New England Biolabs 03 Oct 2014

Can be decreased to 5-15 minutes by using a [Time-Saver™ Qualified enzyme](#).

New England Biolabs 06 Oct 2014

See the [NEB Activity/Performance Chart](#) for the incubation temperatures.