



T Cell Activation with anti-CD3 Antibodies Protocol - Human [↗](#)

Kelsey Knight¹

¹BioLegend

Sep 13, 2018 [dx.doi.org/10.17504/protocols.io.tkbeksn](https://doi.org/10.17504/protocols.io.tkbeksn)

Version 3

BioLegend

Working



Kelsey Knight

BioLegend



EXTERNAL LINK

<https://www.biolegend.com/protocols/t-cell-activation-with-anti-cd3-antibodies-protocol-human/4271/>

PROTOCOL STATUS

Working

MATERIALS TEXT

- Sterile PBS
- Anti-human CD3 Antibody
 - Clone UCHT1 (LEAF™ format, Cat. No. 300413/300414/300432; Ultra-LEAF™ format, Cat. No. 300437/300438)
 - Clone OKT3 (LEAF™ format, Cat. No. 317303/317304/317315; Ultra-LEAF™ format, Cat. No. 317325/317326)
 - Clone HIT 3a (LEAF™ format, Cat. No. 300313/300314; Ultra-LEAF™ format, Cat. No. 300331/300332)
- Cell culture medium (e.g., RPMI-1640 or IMDM supplemented with 10% FBS and 2mM L-glutamine)
- Sterile single-cell suspension of Ficol-Hypaque-purified peripheral blood mononuclear cells, isolated T cells, or T cell subsets
- 96-well flat-bottom tissue culture plates with lids (e.g., Costar® Cat. No. 3596)

* Soluble forms of LEAF™ purified UCHT1 (1 µg/ml) or LEAF™ purified HIT 3a (0.01 – 0.1 µg/ml) may be used to activate T cells from PBMC cell populations.

- 1 Prepare a 10 µg/ml solution of anti-CD3 (clone UCHT1, OKT3, or HIT3a) in sterile PBS.
- 2 Dispense 50 µl of the antibody solution to each microwell of the 96-well assay plate. For the unstimulated control wells, add 50 µl of sterile PBS.

REAGENT



96-well flat-bottom tissue culture plates with lids
by [Corning](#)
Catalog #: [3596](#)

- 3 Seal plate. Incubate at 37°C for 2 hours or 4°C overnight.

02:00:00

- 4 Aseptically decant antibody solution from the microwell plate.

- 5 Wash plate microwells 3 times with sterile PBS. Discard liquid.

- 6 Prepare single cell suspension of cells of interest in supplemented cell culture medium to $1-2 \times 10^6/\text{ml}$.
- 7 Aliquot 200 μl cell suspension into plate microwells. Cover with lid. Incubate at 37°C in 5% CO_2 and 100% humidity for 3 days.



This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited