

# FASP Kit Protocol-ORNL Developed for Bacteriophage

Nathan VerBerkmoes

## Abstract

Collaborator, Nathan VerBerkmoes (Oak Ridge National Labs), worked with Tucson Marine Phage Lab to develop more sensitive proteomics assays for viral isolates which we are now also using for environmental viral concentrates. A new sample prep method (FASP) was optimized to maximize the signal from our commonly low-protein containing samples, and then run using 2d-LC-MS-MS to maximize detection across the large dynamic range in isolate and environmental samples.

**Citation:** Nathan VerBerkmoes FASP Kit Protocol-ORNL Developed for Bacteriophage. **protocols.io**

dx.doi.org/10.17504/protocols.io.ddn25d

**Published:** 21 Jan 2016

## Guidelines

This method was optimized for bacteriophage by Kristen Carrier undergraduate research assistant at Oak Ridge National Laboratory (ORNL), Chemical Science Division using the commercial method from Protein Discovery (Knoxville, TN) as a starting point.

This method was written by Kristen Carrier using the commercial method from Protein Discovery (Knoxville, TN) as a starting point.

Dr. Nathan VerBerkmoes (ORNL, Chemical Science Division) oversaw the optimization process and creation of final method.

This method is designed to prepare small quantities (~1-10µg) of isolate bacteriophage or mixed bacterial/bacteriophage communities for shotgun proteomics via 2d-LC-MS/MS.

Please note this method is not effective for large quantities of material (>100µg), standard solution digest should be used for such samples. The minimum protein amount needed is estimated to be 500ng but this is highly sample dependent.

The main optimization of this method was to simplify it and greatly reduce the time required. This current method takes roughly half the amount of time (actually bench time) as the original commercial method, furthermore its now very straightforward to prepare 8, 16, or 24 samples at a time.

## Protocol

### Day 1

#### Step 1.

Prepare Urea/SDS Solution.

## **PROTOCOL**

### **. Urea/SDS Solution**

CONTACT: [VERVE Team](#)

#### **Step 1.1.**

In falcon tube, add 1 fleck DTT (10mM) to 5mL Tris HCl (provided with FASP kit) or Tris CaCl<sub>2</sub>. Vortex briefly.

#### **Step 1.2.**

Add 1mL Tris buffer + DTT to 1 tube urea (75 µg, provided with FASP kit). Vortex until all powder dissolves.

#### **Step 1.3.**

Combine 300µL Urea solution and 150µL sample (sample can be in any form, mixture of viral proteins and bacteria, ionic or non-ionic detergents, buffers and/or CsCl).

#### **Step 1.4.**

Rock at room temperature for 30 min to 1 hr to lyse bacteria cells/phage particles.

 **DURATION**

01:00:00

Day 1

#### **Step 2.**

Transfer Urea solution + sample to spin filter. Centrifuge at 14,000 x g for 15 min.

 **DURATION**

00:15:00

Day 1

#### **Step 3.**

Add 200µL fresh Urea solution (**no DTT, no SDS**). Centrifuge at 14,000 x g for 15 min.

 **AMOUNT**

200 µl Additional info:

 **DURATION**

00:15:00

## **PROTOCOL**

### **. Urea Solution**

CONTACT: [VERVE Team](#)

#### **Step 3.1.**

1mL Tris CaCl<sub>2</sub> or Tris HCl added to one tube of urea. Vortex until all powder dissolves.

Day 1

#### **Step 4.**

Discard flow-through.

Day 1

#### **Step 5.**

Add 10µL iodoacetamide solution and 90µL Urea solution (no DTT, no SDS). Vortex for 1 min, then incubate without mixing for 20 min in the dark.

 **DURATION**

00:20:00

## [PROTOCOL](#)

### . [Iodoacetamide Solution](#)

CONTACT: [VERVE Team](#)

#### **Step 5.1.**

100µL Urea solution (no DTT, no SDS) added to 1 tube iodoacetamide (provided with FASP kit).  
Pipette up and down 10-15 times to mix well and dissolve.

Day 1

#### **Step 6.**

Centrifuge at 14,000 x g for 10 min.

 [DURATION](#)

00:10:00

Day 1

#### **Step 7.**

Add 100µL Urea solution (no DTT, no SDS). Centrifuge at 14,000 x g for 15 min. Repeat this step twice.

 [DURATION](#)

00:15:00

Day 1

#### **Step 8.**

Discard flow-through.

Day 1

#### **Step 9.**

Add 100µL 50mM ammonium bicarbonate solution (provided with FASP kit). Centrifuge at 14,000 x g for 15 min.

 [DURATION](#)

00:15:00

Day 1

#### **Step 10.**

Transfer filter to new collection tube.

Day 1

#### **Step 11.**

Add 75µL digestion solution. Vortex briefly. Incubate at 37°C for 4 - 18 hours (NO ROCKING)

 [AMOUNT](#)

75 µl Additional info:

 [DURATION](#)

18:00:00

## [PROTOCOL](#)

### . [Digestion Solution](#)

CONTACT: [VERVE Team](#)

#### **Step 11.1.**

75uL ammonium bicarbonate solution added to 20µg trypsin (1 tube).

Day 2

#### **Step 12.**

Add 40µL 50mM ammonium bicarbonate solution. Centrifuge at 14,000 x g for 10 min. Repeat this step once.

 **DURATION**

00:10:00

Day 2

### Step 13.

Add 50µL 0.5 M sodium chloride solution (provided with FASP kit). Centrifuge at 14,000 x g for 10 min.

 **DURATION**

00:10:00

Day 2

### Step 14.

Filtrate contains digested proteins. Add 170µL\* H<sub>2</sub>O + formic acid. Aliquot (150µL x 2)\*, freeze

 **NOTES**

**VERVE Team** 13 Aug 2015

\*Volumes may be adjusted. Final filtrate volume = 130µL

\*Generally two replicates can be obtained from one filter

\*Sample is now ready for 2d-LC-MS/MS, remember a on-line desalting step is needed before SCX salt pulses start.