



Rapid Resistance Testing (RRT) protocol for Rapid Detection of Resistance to Carbapenems and Cephalosporins in Enterobacteriaceae Using Liquid Chromatography Tandem Mass Spectrometry



Version 2

PLOS One

Manal Tadros¹, Alexander Romaschin²

¹Department of Laboratory Medicine & Pathobiology, University of Toronto, Toronto, ON, Canada,

²Division of Microbiology, St. Michael's Hospital, Toronto, ON, Canada

[dx.doi.org/10.17504/protocols.io.uz9ex96](https://doi.org/10.17504/protocols.io.uz9ex96)

Manal Tadros

ABSTRACT

Carbapenemase producing *Enterobacteriaceae* (CPE) are becoming a global healthcare concern. Current laboratory methods for the detection of CPE include screening followed by confirmatory phenotypic and genotypic tests. These processes would generally take ≥ 72 hours, which could negatively impact patient care and Infection Control practices. To this end, we developed a protocol for rapid resistance testing (RRT) to detect hydrolysis in a panel of beta lactam antibiotics consisting of ampicillin, cefazolin, cefotaxime and imipenem, using liquid chromatography tandem mass spectrometry. Ninety-nine beta lactamase producing *Enterobacteriaceae* isolates were used to evaluate the RRT method, 54 isolates were CPE and 45 isolates were Class A or AmpC beta lactamase producing *Enterobacteriaceae* but not carbapenemase producers. We also tested 10 *E. coli* isolates that were susceptible to ampicillin, cefazolin, cefotaxime and imipenem. Receiver Operating Characteristic (ROC) Curves analysis showed that imipenem had a sensitivity and a specificity of 100% for carbapenemase detection at hydrolysis cut off values that are greater than 50% and less than or equal to 80%. The RRT protocol can be conducted in a time frame of less than 2 hours. This preliminary study shows that the rapid resistance testing protocol might have utility for the rapid detection of CPE. Additional work with a greater number and a variety of beta-lactamase producing *Enterobacteriaceae* isolates is required to validate these preliminary findings.

EXTERNAL LINK

<https://doi.org/10.1371/journal.pone.0206842>

THIS PROTOCOL ACCOMPANIES THE FOLLOWING PUBLICATION

Tadros M, Goneau L, Romaschin A, Jarvis M, Matukas L (2018) Rapid detection of resistance to carbapenems and cephalosporins in *Enterobacteriaceae* using liquid chromatography tandem mass spectrometry. PLoS ONE 13(11): e0206842. doi: [10.1371/journal.pone.0206842](https://doi.org/10.1371/journal.pone.0206842)

RRT detailed protocol for
PLOS ONE Final July
30.docx

PROTOCOL STATUS

Working

1

This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited