PNGase F Protocol, Non-Denaturing Conditions

New England Biolabs

Abstract

This is a generic PNGase F protocol with non-denaturing reaction conditions. It is appropriate for both <u>P0704</u> and <u>P0708</u>. Typical reaction conditions are below.

Citation: New England Biolabs PNGase F Protocol, Non-Denaturing Conditions. protocols.io

dx.doi.org/10.17504/protocols.io.cqivud

Published: 31 Jan 2015

Guidelines

When deglycosylating a native glycoprotein it is recommended that an aliquot of the glycoprotein is subjected to the <u>denaturing protocol</u> to provide a positive control for the fully deglycosylated protein. The non-denatured reaction can then be compared to the denatured reaction to determine the extent of reaction completion.

Before start

When deglycosylating a native glycoprotein it is recommended that an aliquot of the glycoprotein is subjected to the <u>denaturing protocol</u> to provide a positive control for the fully deglycosylated protein. The non-denatured reaction can then be compared to the denatured reaction to determine the extent of reaction completion.

Protocol

Step 1.

Combine the following (for a 20 µl total reaction volume):

₽ PROTOCOL

PNGase F Non-Denaturing Mixture

CONTACT: New England Biolabs

NOTES

New England Biolabs 27 Jan 2015

Reactions may be scaled-up linearly to accommodate larger amounts of glycoprotein and larger reaction volumes.

Step 1.1.

Glycoprotein, 1-20 µg

Step 1.2.

10X G7 Reaction Buffer, 2 μl

■ AMOUNT

2 μl Additional info:

Step 1.3.

H2O (if necessary) to make a 20 μl total reaction volume.

Step 2.

Add 2-5 µl PNGase F

NOTES

New England Biolabs 27 Jan 2015

For unit conversion between different suppliers, please reference the Glycobiology Unit Conversion Chart page.

New England Biolabs 27 Jan 2015

If using P0704/P0708, we recommend limiting PNGase F to 1/10 (or less) of the total reaction volume to keep the final glycerol concentration equal to (or less than) 5%.

Step 3.

Mix gently

Step 4.

Incubate reaction at 37°C for 4 - 24 hours

NOTES

New England Biolabs 27 Jan 2015

Optimal incubation times may vary for particular substrates.

New England Biolabs 27 Jan 2015

To deglycosylate a native glycoprotein, longer incubation time as well as more enzyme may be required.

Step 5.

Analyze by method of choice

NOTES

New England Biolabs 27 Jan 2015

The simplest method of assessing the extent of deglycosylation is by mobility shifts on SDS-PAGE gels.