



Aug 23,
2019

Epigenomic profiling of neuroblastoma cell lines

Kristen Upton¹, Robyn T. Sussman¹, Khushbu Patel², Gregory P. Way³, Rebecca N. Adams⁴, Gregory I. Sacks¹, Rebecca N. Adams⁴, Paolo Fortina⁴, John M. Maris^{1,5}, Jo Lynne Rokita^{1,2,6}

¹Division of Oncology, Children's Hospital of Philadelphia, Philadelphia, Pennsylvania 19104, USA,

²Department of Bioinformatics and Health Informatics, Children's Hospital of Philadelphia, Philadelphia, Pennsylvania 19104, USA, ³Genomics and Computational Biology Graduate Group, University of Pennsylvania, Philadelphia, PA, 19104, USA, ⁴Cancer Genomics and Bioinformatics Laboratory, Sidney Kimmel Cancer Center, Philadelphia, Pennsylvania 19107, USA, ⁵Department of Pediatrics, Perelman School of Medicine at the University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA, ⁶Center for Data-Driven Discovery in Biomedicine, Children's Hospital of Philadelphia, Philadelphia, Pennsylvania 19104, USA

2

Works for me

[dx.doi.org/10.17504/protocols.io.6r3hd8n](https://doi.org/10.17504/protocols.io.6r3hd8n)



Kristen Upton



ABSTRACT

This protocol explains the process of how we collected MYCN, MYC, and Histone ChIP-Seq data, as well as ATAC-Seq data for neuroblastoma cell lines. This protocol is comprised of three sections: Cell Growth and Expansion for care of neuroblastoma cell lines, ChIP-Seq protocol, and ATAC-Seq Protocol. Table 1, within the document, outlines which data was collected for each cell line.

□

ChIP-Seq-ATAC-Seq-
Protocol.pdf

1



This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited