

Sanger sequencing protocol - BRCA1/2 mutations are not a common cause of malignant melanoma in the Polish population 👄

PLOS One

Oct 05, 2018

Tadeusz Dębniak1¹, Rodney J Scott², Bohdan Górski¹, Bartłomiej Masojć³, Andrzej Kram¹, Romuald Maleszka¹, Cezary Cybulski¹, Katarzyna Paszkowska-Szczur¹, Aniruddh Kashyap¹, Dawid Murawa⁴, Karolina Malińska¹, Magdalena Kiedrowicz¹, Emilia Rogoża-Janiszewska¹, Helena Rudnicka¹, Jakub Deptuła¹, Paweł Domagała¹, Wojciech Kluźniak¹, Marcin R. Lener¹, Jan Lubiński¹

¹Pomeranian Medical University in Szczecin, ²University of Newcastle and the Hunter Medical Research Institute, ³West Pomeranian Oncology Cente, ⁴Greater Poland Cancer Center

dx.doi.org/10.17504/protocols.io.ta7eihn



Bartlomiej Masojc 🚱



ABSTRACT

In the study we have used a standard Thermo Fisher Scientific Sanger sequencing protocol which can be found under this link:

http://www.ramaciotti.unsw.edu.au/wp-content/uploads/2016/08/sequencing_handbook_FLR.pdf

EXTERNALLINK

https://doi.org/10.1371/journal.pone.0204768

THIS DOCUMENT ACCOMPANIES THE FOLLOWING PUBLICATION

Dębniak T, Scott RJ, Górski B, Masojć B, Kram A, Maleszka R, Cybulski C, Paszkowska-Szczur K, Kashyap A, Murawa D, Malińska K, Kiedrowicz M, Rogoża-Janiszewska E, Rudnicka H, Deptuła J, Domagała P, Kluźniak W, Lener MR, Lubiński J (2018) BRCA1/2 mutations are not a common cause of malignant melanoma in the Polish population. PLoS ONE 13(10): e0204768. doi: 10.1371/journal.pone.0204768

Document body

This is an open access document distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited