

Jul 12, 2019

SPARC Serotonin 2B Receptor (5-HT2BR) Immunohistochemistry Protocol in Rat Tissues Labeled with Cholera Toxin B-fragment

Elisa Gonzalez-Rothi¹, Yasin Seven¹, Latoya Allen¹, Marissa Ciesla¹, Gordon Mitchell¹ University of Florida

1 Works for me dx.doi.org/10.17504/protocols.io.2kigcue



ABSTRACT

This protocol describes the immunofluorescent labeling technique used to identify serotonin 2B receptor expression in CtB-labelled phrenic motor neurons and within a defined region of interest surrounding phrenic motor neurons.

- Day 1: primary antibodies required: 5-HT2BR: Mouse anti-5-HT2BR (SantaCruz Biotechnology #C-6; SC-376878) Cholera toxin B-fragment: Goat anti-CT-B (Millipore #227040)
- 2 Place 40um transverse spinal cord sections into 1xPBS in 12 well cell culture plates
- 3 5x washes in 1xPBS for 5 minutes each at room temperature
- 4 Blocking: place tissues in 5% Normal Horse Serum (NHS) in 1xPBS-Triton (0.1%) +0.5% bovine serum albumin (BSA) for 60 minutes at room temperature
- 5 Primary Antibody Incubation: Incubate tissues in: 5%NHS in 1xPBS-Triton (0.1%), Mouse anti-5-HT2BR (1:1000), and Goat anti-Ct-B (1:2500) overnight at 4 degrees C
- 6 Day 2: secondary antibodies required: AlexaFluor 488: donkey anti-goat (Invitrogen, Ref#A11055) AlexaFluor 594: donkey anti-mouse (Invitrogen, Ref#A21203)
- 7 5x washes in 1xPBS for 5 minutes each at room temperature
- 8 Secondary Antibody Incubation: Incubate tissues in: 5%NHS in 1xPBS-Triton (0.1%), donkey anti-goat (1:1000), and donkey anti-mouse (1:1000) for 2 hours at room temperature
- 9 5x washes in 1xPBS for 5 minutes each at room temperature
- 10 Mount tissues on Superfrost Plus microscope slides
- 11 Allow slides to dry overnight

This is an open access protocol distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited