



Jul 16, 2019

SPARC_Duke_Grill_OT2-OD025340_RatVagusNerveCollectionHistologyMicroscopy

J. Ashley Ezzell¹, Nicole A. Pelot², Kara A. Clissold¹, Warren M. Grill²¹University of North Carolina, ²Duke University

1

Works for me

dx.doi.org/10.17504/protocols.io.ww3ffgn



Nikki Pelot



ABSTRACT

Protocol for collection, histological processing, and imaging of rat vagus nerves.

MATERIALS TEXT

- Adult Sprague-Dawley rats from Charles River
- 4% paraformaldehyde
- Mordant & tissue dye
- Ethanol
- Clearite
- Paraffin
- Xylene
- Bouin's fixative (mordant)
- Weigert's iron hematoxylin solution
- Biebrich scarlet-acid fuchsin solution
- Phosphomolybdic-phosphotungstic acid solution
- Aniline blue solution
- Acetic acid
- Microscope with color camera

Collect rat vagus nerve samples.

- 1 We collected vagus nerve samples from perfused Sprague-Dawley rats that were used in other experiments approved by the Duke University Institutional Animal Care and Use Committee.
- 2 We perfused the rats with ~300 mL phosphate buffered saline or until the liver was cleared, followed by ~300 mL 4% paraformaldehyde or until the joints and tissues were stiff.
- 3 We collected samples of the carotid sheath in the neck bilaterally, where the vagus nerve courses straight with the common carotid artery, from the clavicle up to where the nerve and carotid intermingle with non-parallel other nerves. We measured from the "valley" of the common carotid bifurcation to the center of each sample.
- 4 We collected the esophagus from the diaphragm to the stomach with the subdiaphragmatic vagus nerves attached.
- 5 We dyed the rostral end of each sample green to maintain orientation during processing.
- 6 We placed each sample between two histology sponges in a mega-sized histology cassette. We placed the cassettes in a tub with 4% paraformaldehyde in a 4°C refrigerator.

Perform histological processing.

- 7 We rinsed each sample with deionized water.
- 8 We processed each sample on the long cycle in the Leica ASP300S Tissue Processor for ~10 hours: 70, 80, 95, 95, 100, 100, 100% ethanol for 30, 35, 40, 40, 40, 40, 40 min, respectively; Clearite for 50 min, three times; paraffin wax for 50 min, three times.
- 9 We embedded each sample in a paraffin block oriented to obtain transverse sections.
- 10 After trimming to the center of the sample, we collected 5 µm sections, placing two serial sections per microscope slide for fifteen slides.
- 11 The slides were air dried overnight at room temperature, then baked at 37°C overnight.
- 12 Of the 15 slides per sample, we stained slides 2 and 14 with Masson's trichrome as follows.
- 13 The slides were baked at 60°C for 1.5 hours and then cooled overnight.
- 14 We deparaffinized the slides and hydrated them to distilled water: xylene (2x 6 min), 100% ethanol (5 min), 95% ethanol (4 min), 70% ethanol (3 min), dH₂O (2x 1 min).
- 15 We placed the slides in Bouin's fixative (mordant) at room temperature overnight.
- 16 We washed the slides in running tap water until the yellow color disappeared (~10 minutes).
- 17 We rinsed the slides in distilled water.
- 18 We placed the slides in Weigert's Iron Hematoxylin solution for 10 min.
- 19 We washed the slides in running tap water for 10 min.
- 20 We placed the slides in Biebrich Scarlet-Acid Fuchsin solution for 5 min.
- 21 We washed the slides in running tap water for 2 min.
- 22 We placed the slides in Phosphomolybdic-Phosphotungstic Acid solution for 10 min.

- 23 We transferred the slides directly to Aniline Blue solution for 3 min.
- 24 We differentiated the counterstain (A. blue) in 1% Acetic Acid solution for 1 min.
- 25 We dehydrated, cleared, and coverslipped the slides: 95% ethanol (1 min), 100% ethanol (4 min), xylene (5 min).

Perform microscopy.

- 26 Each sample was imaged at 20x using a Nikon Ti2 microscope with a Photometrics Prime 95B-25MM camera (Nikon Instruments Inc.). We selected the best of four slices for each sample based on the quality of the slice (no tearing or fraying).



This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited