

SPARC Serotonin (5-HT) Immunohistochemistry Protocol in Rat Tissues Labeled with Cholera **Toxin B-fragment**

Elisa Gonzalez-Rothi¹, Yasin Seven¹, Latoya Allen¹, Marissa Ciesla¹, Gordon Mitchell¹ ¹University of Florida

1 Works for me dx.doi.org/10.17504/protocols.io.2kggctw



ABSTRACT

This protocol describes the immunofluorescent labeling technique used to identify serotonin expression in CtB-labelled phrenic motor neurons and within a defined region of interest surrounding phrenic motor neurons.

- Day 1: primary antibodies required: 5-HT: Rabbit anti-5-HT (Immunostar #20080) Cholera toxin B-fragment: Goat anti-CT-B (Millipore #227040)
- Place 40um transverse spinal cord sections into 1xPBS-Triton (0.1%) in 12 well cell culture plates
- 5x washes in 1xPBS-Triton (0.1%) for 5 minutes each at room temperature 3
- Antigen retrieval: place tissues into Heat-Induced Epitope Retrieval (TissuePro, Cat#: HIER01-32R) for 30 minutes at 85 degrees C
- 5x washes in 1xPBS-Triton (0.1%) for 5 minutes each at room temperature
- Blocking: place tissues in 5% Normal Donkey Serum (NDS) in 1xPBS-Triton (0.1%) for 60 minutes at room temperature
- Primary Antibody Incubation: Incubate tissues in: 5% NDS in 1xPBS-Triton (0.1%), Rabbit anti-5-HT (1:2000), and Goat anti-Ct-B (1:2500) for 1 hour at room temperature
- Continue primary antibody incubation overnight at 4 degrees C
- Day 2: secondary antibodies required: AlexaFluor 488: donkey anti-goat (Invitrogen, Ref#A11055) AlexaFluor 594: donkey anti-rabbit (Invitrogen, Ref#A21207)
- 5x washes in 1xPBS-Triton (0.1%) for 5 minutes each at room temperature 10
- Secondary Antibody Incubation: Incubate tissues in: 5% NDS in 1xPBS-Triton (0.1%), donkey anti-goat (1:1000), and donkey anti-rabbit 11

(1:500) for 2 hours at room temperature

- 12 5x washes in 1xPBS for 5 minutes each at room temperature
- 13 Mount tissues on Superfrost Plus microscope slides
- 14 Allow slides to dry overnight
- 15 Coverslip with VectaShield Antifade Hard Set Mounting Medium (Cat#:H-1400)

This is an open access protocol distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited