

Jun 19 2019

Transformation of *Diplonema papillatum* using V5+Neo^R construct

Binnypreet Kaur¹, Drahomíra Faktorová², Julius Lukes²

¹1 Institute of Parasitology, Biology Centre, Czech Academy of Sciences 2 Faculty of Sciences, University of South Bohemia, České Budějovice (Budweis), Czech Republic, ²Research Scientist

Working dx.doi.org/10.17504/protocols.io.4digs4e

Protist Research to Optimize Tools in Genetics (PROT-G) Julius Lukes





ABSTRACT

p57-V5+Neo^R plasmid contains V5-tagged aminoglycoside 3'-phosphotransferase gene (APT; conferring resistance to neomycin) flanked by partial regulatory sequences derived from the hexokinase gene of the kinetoplastid Blastocrithidia (strain p57).

Plasmid GenBank accession number = MN047315

MATERIALS TEXT

p57-V5+NeoR plasmid.jpg

- p57-V5+Neo^R plasmid contains V5-tagged aminoglycoside 3'-phosphotransferase gene (APT; conferring resistance to neomycin) flanked by partial regulatory sequences derived from the hexokinase gene of the kinetoplastid Blastocrithidia (strain p57).
- This 1779 bp long sequence is surrounded by Swal restriction sides and was cloned in the pBluescript II SK(+) backbone.
- Before transformation, the p57-V5+ Neo^R plasmid was cut using Swal, run on the gel and a fragment of 1779 bp containing p57-V5+ Neo^R construct was gel purified
- 3 ug of purified construct was electroporated into D. papillatum cells using an established protocol (Kaur et al,. 2018). About 18 h after electroporation, a scale of 45 - 83 ug/ml G418 (Geneticin) was added to the media and after about 2 weeks G418-resistant clones were recovered.

This is an open access protocol distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited