

Euplotes crassus transformation using FuGene HD Transfection Reagent as vehicle

RACHELE CESARONI

Abstract

Citation: RACHELE CESARONI Euplotes crassus transformation using FuGene HD Transfection Reagent as vehicle. **protocols.io**

dx.doi.org/10.17504/protocols.io.g3ebyje

Published: 26 Jan 2017

Protocol

Step 1.

2×10^4 well-fed cells (we used E.coli as the only food source) were collected by centrifugation (400 rcf for 3 minutes).

Step 2.

Cells were washed twice with artificial sea water (see attachment for the recipe) and once with 500 mM sorbitol, 0.5 mM Tris-HCl pH 7.0 (400 rcf for 3 minutes each time), where they were also resuspended in 50 μ l (to get this small volume it may be required an additional minute of centrifugation).

📌 NOTES

Rachele Cesaroni 26 Jan 2017

Recipe for complete sea water (1 L):

36 g Reef Crystals

1 ml Walne's solution

1 ml of 10 μ g/ml FeSO_4

0.2 ml of 2 mg/ml thiamine (light sensitive; store at 4°C)

Add distilled water to 1 L

Step 3.

Cells were transferred in 96-well plates for transfection.

■ ANNOTATIONS

Estienne Swart 27 Jan 2017

How many wells were used?

Rachele Cesaroni 30 Jan 2017

Two wells: one for the negative control and one for the cells with FuGene/DNA complexes.

Step 4.

FuGene HD Transfection Reagent was allowed to reach room temperature and mixed by inverting or vortexing briefly (if a precipitate is visible, briefly warm at 37°C and then let it reach room temperature).

Step 5.

90-98 µl of medium at room temperature (500 mM sorbitol, 0.5 mM Tris-HCl pH 7.0) and 4 µg of DNA in MilliQ H₂O (0.2-1 µg/µL) were added to an Eppendorf tube and vortexed (after adding the DNA the final volume must be 100 µl).

Step 6.

6 µl of FuGene HD Transfection Reagent were added directly to the medium and mixed immediately.

Step 7.

FuGene HD Transfection Reagent and DNA mixture were incubated for 15 minutes at room temperature.

Step 8.

10 µl of the FuGene HD Transfection reagent and DNA mixture were added to the wells containing 50 µl of cells in medium (everything was mixed by pipetting).

Step 9.

One hour after addition of FuGene/DNA complexes, 50 µl of artificial sea water were added to the cells.

Step 10.

After another hour other 50 µl of artificial sea water were added to the cells.

Step 11.

An hour later cells were harvested (400 rcf for 3 minutes) and washed twice with artificial sea water (400 rcf for 3 min each time),

Step 12.

Cells were transferred in depression wells.

Step 13.

Cells were incubated at least for three days at 24°C, then analyzed.