



## Adult fecundity [↗](#)

PLOS Genetics

Wen Aw<sup>1</sup>

<sup>1</sup>z3314717@unsw.edu.au

[dx.doi.org/10.17504/protocols.io.rq2d5ye](https://doi.org/10.17504/protocols.io.rq2d5ye)

Cage Studies

Wen Aw

### EXTERNAL LINK

<https://doi.org/10.1371/journal.pgen.1007735>

### THIS PROTOCOL ACCOMPANIES THE FOLLOWING PUBLICATION

Aw WC, Towarnicki SG, Melvin RG, Youngson NA, Garvin MR, Hu Y, Nielsen S, Thomas T, Pickford R, Bustamante S, Vila-Sanjurjo A, Smyth GK, Ballard JWO (2018) Genotype to phenotype: Diet-by-mitochondrial DNA haplotype interactions drive metabolic flexibility and organismal fitness. PLoS Genet 14(11): e1007735. doi: [10.1371/journal.pgen.1007735](https://doi.org/10.1371/journal.pgen.1007735)

### PROTOCOL STATUS

**Working**

- 1 Female fecundity was assayed using flies that had been transferred from instant food to each experimental diet for one generation.
- 2 Briefly, 47 female flies of each genotype and diet, ranging from 3-5 days old were randomly transferred into separate 6 mL glass vials (Sigma-Aldrich) and allowed to oviposit for 24 hours.
- 3 The numbers of eggs in each vial were then counted under the microscope.



This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited