

Neutrophil-lymphocyte ratio predicts the outcome of intracerebral hemorrhage: A meta-analysis and systematic review

weiduan zhuang

Abstract

Background: The neutrophil-lymphocyte ratio (NLR) is increasingly known as an indicator of systemic inflammation. It is used as a predictor for clinical outcomes in cancers and cardiovascular disease. However, its relationship with intracerebral hemorrhage (ICH) is still disputed.

Aims: We sought to evaluate the association between NLR and ICH in this meta-analysis.

Methods: Potentially relevant articles were searched in PubMed, Cochrane Library, Medline and EMBASE until April 8, 2018, with English restriction. Efficacy outcomes included major disability at 90 days, short-term mortality or in-hospital mortality. Odds ratios (ORs) with 95% confidence intervals (95% CIs) were pooled to estimate the prognostic role of NLR in ICH.

Results: A total of 7 trials with 2176 patients were included. The results of our meta-analysis revealed that high NLR had a higher risk of major disability at 90 days (OR: 2.20; 95% CI: 1.27-3.81) and higher mortality at short-term (OR: 1.31; 95% CI: 1.02-1.68); without statistically significant association with in-hospital mortality (OR: 1.02; 95% CI: 0.91-1.15).

Conclusions: Our meta-analysis proves that higher NLR was a predictor of major disability at 90 days and higher mortality at short term in patients with ICH, but not a predictor of in-hospital mortality.

Citation: weiduan zhuang Neutrophil-lymphocyte ratio predicts the outcome of intracerebral hemorrhage: A meta-analysis and systematic review. **protocols.io**

dx.doi.org/10.17504/protocols.io.qzwdx7e

Published: 15 Jun 2018

Protocol

Background

Step 1.

The neutrophil-lymphocyte ratio (NLR) is increasingly known as an indicator of systemic inflammation. It is used as a predictor for clinical outcomes in cancers and cardiovascular disease. However, its relationship with intracerebral hemorrhage (ICH) is still disputed.

Aims

Step 2.

We sought to evaluate the association between NLR and ICH in this meta-analysis.

Methods

Step 3.

Potentially relevant articles were searched in PubMed, Cochrane Library, Medline and EMBASE until April 8, 2018, with English restriction. Efficacy outcomes included major disability at 90 days, short-term mortality or in-hospital mortality. Odds ratios (ORs) with 95% confidence intervals (95% CIs) were pooled to estimate

the prognostic role of NLR in ICH.

Results

Step 4.

A total of 7 trials with 2176 patients were included. The results of our meta-analysis revealed that high NLR had a higher risk of major disability at 90 days (OR: 2.20; 95% CI: 1.27-3.81) and higher mortality at short-term (OR: 1.31; 95% CI: 1.02-1.68); without statistically significant association with in-hospital mortality (OR: 1.02; 95% CI: 0.91-1.15).

Conclusions

Step 5.

Our meta-analysis proves that higher NLR was a predictor of major disability at 90 days and higher mortality at short term in patients with ICH, but not a predictor of in-hospital mortality.
