



Mar 28,  
2019

Working

## Determination of flavonoid content [↗](#)

PLOS One

Jorge Carlos Ruiz Ruiz<sup>1</sup>

<sup>1</sup>Universidad Anáhuac Mayab

[dx.doi.org/10.17504/protocols.io.sfnebme](https://doi.org/10.17504/protocols.io.sfnebme)

Jorge Carlos Ruiz Ruiz

### ABSTRACT

Flavonoid content determination using the aluminum chloride method.

### EXTERNAL LINK

<https://doi.org/10.1371/journal.pone.0213493>

### THIS PROTOCOL ACCOMPANIES THE FOLLOWING PUBLICATION

Dewanto V, Wu X, Adom KK, Liu RH. Thermal processing enhances the nutritional value of tomatoes by increasing total antioxidant activity. J Agric Food Chem. 2002; 50: 3010-3014.

### PROTOCOL STATUS

**Working**

### MATERIALS

NAME	CATALOG #	VENDOR
Sodium nitrite	<a href="#">View</a>	P212121
water		
aluminum chloride		<a href="#">Sigma Aldrich</a>
sodium carbonates		<a href="#">Sigma Aldrich</a>
catechin		<a href="#">Sigma Aldrich</a>

### SAFETY WARNINGS

- 1 An aliquot of sample (0.1 mL) was mixed with 0.4 ml distilled water in a 1.5 mL microcentrifuge tube.
- 2 0.03 mL of 5% NaNO<sub>2</sub> was added and the mixture was allowed to react for 5 min.
- 3 Following this, 0.03 mL of 10% AlCl<sub>3</sub> was added and the mixture stood for a further 5 min.
- 4 Finally, the reaction mixture was treated with 0.2 mL of 1 M Na<sub>2</sub>CO<sub>3</sub> and 0.24 mL distilled water, and the absorbance at 510 nm.

5 Catechin (Sigma-Aldrich) was used as the standard and the results were expressed as  $\mu\text{g/mL}$  of catechin equivalents (CE).



This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited