



May 09,  
2019

Working

## U Cinn - Non-invasive Measurement of Intestinal Fat Absorption [↗](#)

Patrick Tso<sup>1</sup>, Dana Lee<sup>1</sup>

<sup>1</sup>University of Cincinnati

dx.doi.org/10.17504/protocols.io.xmtfk6n

Mouse Metabolic Phenotyping Centers

Tech. support email: [info@mmpc.org](mailto:info@mmpc.org)

Lili Liang

### ABSTRACT

#### Summary:

Dietary fat containing 5% sucrose polybehenate, a non-absorbable food additive, is fed in a semi-synthetic diet containing absorbable fat to rodents. The animals eat the diet ad lib for 3 days. Two or more fecal pellets are collected on days 3 and 4. Fat absorption is calculated from the ratios of behenic acid to other fatty acids in the diet and in the feces as analyzed by gas chromatography of fatty acid methyl esters.

### EXTERNAL LINK

<https://mmpc.org/shared/document.aspx?id=197&docType=Protocol>

### MATERIALS

NAME	CATALOG #	VENDOR	CAS NUMBER	RRID
Rodent Diet With 45 kcal% Fat and Modification To 16% Fat (Mostly Safflower Oil or Butter Fat) With 5% Fat as Sucrose Polybehenate	D07031501-02	Research Lab Diets		

## 1 Animal Feeding and Fecal Collection:

**MONDAY:** REPLACE CHOW WITH TEST DIET

**TUESDAY:** CONTINUE TEST DIET,  
NEW CAGE BEDDING BEFORE FEEDING PERIOD

**WEDNESDAY:** CONTINUE TEST DIET  
**(DAY 3 – RETURN)**  
COLLECT 5-8 FECAL PELLETS/ANIMAL  
CHANGE CAGE BEDDING BEFORE FEEDING PERIOD

**THURSDAY:** COLLECT 5-8 FECAL PELLETS/ANIMAL  
**(DAY 4 – RETURN)**



This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited