

Jun 15, 2019

Scanning electron microscopy

PLOS One

Sue Lin¹

¹Institute of Life Sciences, Wenzhou University, Wenzhou, China

Working dx.doi.org/10.17504/protocols.io.zz4f78w



EXTERNAL LINK

https://doi.org/10.1371/journal.pone.0218029

THIS PROTOCOL ACCOMPANIES THE FOLLOWING PUBLICATION

Lin S, Miao Y, Su S, Xu J, Jin L, Sun D, Peng R, Huang L, Cao J (2019) Comprehensive analysis of Ogura cytoplasmic male sterility-related genes in turnip (*Brassica rapa* ssp. *rapifera*) using RNA sequencing analysis and bioinformatics. PLoS ONE 14(6): e0218029. doi: 10.1371/journal.pone.0218029

MATERIALS TEXT

2.5% glutaraldehyde in phosphate buffer (pH7.0), phosphate buffer (pH7.0), 1% OsO₄ in phosphate buffer (pH7.0), ethanol (50%, 70%, 80%, 90%, 95% and 100%), iso-amyl acetate, liquid CO₂, gold-palladium.

- Double fixation: The specimen was first fixed with 2.5% glutaraldehyde in phosphate buffer (pH7.0) for more than 4 hours; washed three times in the phosphate buffer; then postfixed with 1% OsO₄ in phosphate buffer (pH7.0) for 1 hour and washed three times in the phosphate buffer.
- Dehydration: The specimen was first dehydrated by a graded series of ethanol (50%, 70%, 80%, 90%, 95% and 100%) for about 15 to 20 minutes at each step, transferred to the mixture of alcohol and iso-amyl acetate (v:v=1:1) for about 30 minutes, then transferred to pure iso-amyl acetate for about 1 hour. In the end, the specimen was dehydrated in Hitachi Model HCP-2 critical point dryer with liquid CO₂.
- 3 Coating and observation: The dehydrated specimen was coated with gold-palladium in an Eiko Model IB5 ion coater (Eiko Engineering Company, Ibaraki, Japan) and observed in a Hitachi Model TM-1000 scanning electron microscope (Tokyo, Japan).

This is an open access protocol distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited