

# An assessment on a large geographic scale of Eurasian inland saline surface waters

Version 3

Emil Boros, Marina Kolpakova

## Abstract

The major ion concentration data of sodium (Na), potassium (K), calcium (Ca), magnesium (Mg), chloride (Cl), sulphate (SO<sub>4</sub>), bicarbonate (HCO<sub>3</sub>), carbonate (CO<sub>3</sub>) and pH (if it was coupled with ion data) were input into the database. The data were drawn from a large geographic scale across Eurasia (Austria, China, Hungary, Kazakhstan, Mongolia, Russia, Serbia, Turkey) and from a large number of saline lakes and pans (N=220) with minimum a 1.0 g L<sup>-1</sup> salinity threshold. The 1.0 g L<sup>-1</sup> salinity threshold was selected based on a former study (Boros et al., 2014), where this threshold was experimentally found to be the characteristic boundary of soda ecosystems. Salinity was estimated by the sum of measured concentrations of eight major ions (Na, K, Ca, Mg, Cl, SO<sub>4</sub>, HCO<sub>3</sub>, CO<sub>3</sub>). As generally known, sodium is by far the most common cation in saline lakes and necessarily the dominant cation in soda type lakes. Therefore, sites were excluded from the dataset if Na was not the most abundant ion. If seasonal or annual water data were available, mean values were put into database. Most of the data came from papers (sources are indicated in the table).

**Citation:** Emil Boros, Marina Kolpakova An assessment on a large geographic scale of Eurasian inland saline surface waters. **protocols.io**

[dx.doi.org/10.17504/protocols.io.rikd4cw](https://dx.doi.org/10.17504/protocols.io.rikd4cw)

**Published:** 06 Jul 2018

## Protocol

### Chemical assessment of Eurasian inland saline surface waters

#### Step 1.

##### Data collection and measurements

Major ion concentration data including sodium (Na<sup>+</sup>), potassium (K<sup>+</sup>), calcium (Ca<sup>2+</sup>), magnesium (Mg<sup>2+</sup>), chloride (Cl<sup>-</sup>), sulfate (SO<sub>4</sub><sup>2-</sup>), bicarbonate (HCO<sub>3</sub><sup>-</sup>), carbonate (CO<sub>3</sub><sup>2-</sup>), and pH (if it was coupled with ion data) were compiled into a database. The data were drawn from a large geographic scale across Eurasia (Austria, China, Hungary, Kazakhstan, Mongolia, Russia, Serbia, and Turkey) and from a large number of saline lakes and pans (N = 220) with a minimum of 1.0 g L<sup>-1</sup> salinity threshold. The 1.0 g L<sup>-1</sup> salinity threshold was selected on the basis of a former study [27], where this threshold was experimentally found to be the characteristic boundary of soda ecosystems. Salinity (TDS) was estimated by the sum of major ion concentrations (Na<sup>+</sup>, K<sup>+</sup>, Ca<sup>2+</sup>, Mg<sup>2+</sup>, Cl<sup>-</sup>, SO<sub>4</sub><sup>2-</sup>, HCO<sub>3</sub><sup>-</sup>, and CO<sub>3</sub><sup>2-</sup>) and 1 mg L<sup>-1</sup> accuracy was used for all input concentration data independently of the theoretical accuracy of original measurements, therefore decimals were rounded to integer value. As generally known, sodium is by far the most common cation in saline lakes and necessarily the dominant cation in soda type of lakes [1]. Therefore, sites with the most abundant ion other than Na<sup>+</sup> were excluded from the dataset. If seasonal or annual water data were available, we used mean values for the database. Most of the data came from papers [1,15,26-39] published in the period from 1956 to 2017. However, we cannot estimate a reliable universal accuracy and reproducibility for all data in this study, because none of the published papers reviewed provide data on uncertainty, furthermore, several of them were carried out by different kinds of national standard methods of ion concentration measurements, but all of them are based on the same international methods of ICP-

MS for cations (ASTM 3500-Na-C, -K-C, -Ca-C, -Mg-C), alkalinity titration (ASTM 2320 B), chloride argentometric titrimetry (ASTM 4500-ClB) as well as sulfate ions turbidimetry (ASTM 4500-SO<sub>4</sub><sup>2-</sup>E). The collected published data were supplemented with some unpublished sample series in the case of Mongolia and Kazakhstan. They are indicated in the attached table as "Author's unpublished data". The pH was measured on site using a WTW multiline field instrument with a SenTix 41 electrode. The samples were filtered through GF5 glass fiber filters (0.4 µm nominal pore size) in the laboratory. For the determination of cations (Na<sup>+</sup>, K<sup>+</sup>, Ca<sup>2+</sup>, and Mg<sup>2+</sup>), samples were analyzed with the EPA 6020 standard method of inductively coupled plasma mass spectrometry [40] with 2% accuracy. Anion concentrations were measured according to Hungarian standard analytical methods (MSZ). Chloride ion concentration was determined by the argentometric method (MSZ 448-15:1982). Sulfate ion was precipitated in a strong acidic medium with barium chloride, and the resulting turbidity was measured photometrically at 405 nm and compared with standard solutions (MSZ 12750-16:1998). Alkalinity was determined by titration, and the concentration of HCO<sub>3</sub><sup>-</sup>, CO<sub>3</sub><sup>2-</sup>, and OH<sup>-</sup> ions was calculated using the Hungarian standard MSZ 448-11:1986. All of anions and alkalinity were measured with 5% accuracy.

## References

- [1] Hammer UT. Saline lake ecosystems of the World. Dr W. Junk Publishers, The Hague; 1986.
- [15] Shvartsev L, Kolpakova MN, Isupov VP, Vladimirov AG, Ariunbileg S. Geochemistry and Chemical Evolution of Saline Lakes of Western Mongolia. *Geochemistry International*. 2014; 52(5): 388-403.
- [26] Zheng M, Jiayou T, Junying L, Fasheng Z. Chinese saline lakes. *Hydrobiologia*. 1993; 267: 23-36.
- [27] Boros E, Horváth Zs, Wolfram G, Vörös L. Salinity and ionic composition of the shallow astatic soda pans in the Carpathian Basin. *Annales de Limnologie, International Journal of Limnology*. 2014; 50(1): 59-69.
- [28] Wolfram G, Déri L, Zech S, editors. Neusiedler See strategic study – Phase 1. (in Hungarian and German), Fertő tó Stratégiai tanulmány – 1. fázis. Tanulmány a Magyar – Osztrák Vízügyi Bizottság megbízásából. Bécs – Szombathely; 2014.
- [29] Williams WD. Chinese and Mongolian saline lakes: a limnological overview. *Hydrobiologia*. 1991; 210: 39-66.
- [30] Boros E, Jurecska L, Tatár E, Vörös L, Kolpakova M. Chemical composition and trophic state of shallow saline steppe lakes in central Asia (North Kazakhstan). *Environmental Monitoring and Assessment*. 2017; 189: 546, <https://doi.org/10.1007/s10661-017-6242-6>
- [31] Gaskova OL, Strakhovenko VD, Ovdina EA. Composition of brines and mineral zoning of the bottom sediments of soda lakes in the Kulunda steppe (West Siberia). *Russian geology and geophysics*. 2017; 58(10): 1199-1210. doi: 10.1016/j.rgg.2016.09.034
- [32] Williams WD, Aladin NV. The Aral Sea: recent limnological changes and their conservation significance. *Aquatic Conservation: Marine and Freshwater Ecosystems*. 1991; 1: 3–23.
- [33] Doi H, Kikuchi E, Mizota C, Satoh N, Shikano S, Yurlova N, Yadrenkina E, Zuykova E. Carbon, nitrogen, and sulfur isotope changes and hydro-geological processes in a saline lake chain. *Hydrobiologia*. 2004; 529: 225-235.
- [34] Gaskova OL, Kolpakova MN, Naymushina OS, Krivonogov SK. Geochemical processes controlling the water chemistry of saline lakes in the North Kazakhstan Region. *International Multidisciplinary Scientific GeoConference Surveying Geology and Mining Ecology Management*. 2017; 17(31): 325-333.
- [35] Gorlenko VM, Buryukhaev SP, Matyugina EB, Borzenko SV, Namsaraev Z., Bryantseva IA, Boldareva EN, Sorokin DYU, Namsaraev BB. Microbial Communities of the Stratified Soda Lake Doroninskoe (Transbaikal Region). *Microbiology*. 2010; 79(3); 390-401.

- [36] Guseva NV, Kopylova YuG, Hvachevskaya AA, Smetanina IV. The chemical composition of the salt lakes of the North-Minusinsk depression, Khakassia. Bulletin of the Tomsk Polytechnic University. 2012; 321(1): 163–168. (in Russian)
- [37] Kolpakova MN, Borzenko SV, Isupov VP, Shatskaya SS, Shvartsev SL. Hydrochemistry and geochemical classification of salt lakes steppes of the Altai territory. Water chemistry and ecology. 2015; 1: 18–23. (In Russian)
- [38] Zamana LV, Borzenko SV. Hydrochemical regime of saline lakes in the Southeastern Transbaikalia. Geography and Natural Resources. 2010; 31(4): 370–376.
- [39] Kolpakova M.N, Gaskova O.L., Naymushina O.S., Krivonogov S.K. Ebeity Lake, Russia: chemical-organic and mineral composition of water and bottom sediments. Bulletin of the Tomsk Polytechnic University. Geo Assets Engineering. 2018. V. 329. 1. 111–123.
- [40] EPA (U.S. Environmental Protection Agency). "Method 6020. Inductively Coupled Plasma-Mass Spectrometry." Test Methods for Evaluating Solid Wastes: Physical/Chemical Methods, EPA SW-846, Third Ed., Vol. I, Section A, Chapter 3 (Inorganic Analytes); 1994.
-