

Working

U Mass - Glucose Tolerance Test 👄

Jason Kim¹

¹University of Massachusetts

dx.doi.org/10.17504/protocols.io.xxafpie

Mouse Metabolic Phenotyping Centers Tech. support email: info@mmpc.org



Lili Liang 69



ABSTRACT

Summary:

Glucose tolerance test measures systemic clearance of glucose following an intraperitoneal bolus injection of 20% dextrose. This experiment measures insulin sensitivity in awake mice assuming that there are no alterations in the animal's pancreatic β -cell function and insulin secretion. Insulin sensitivity is altered in obese mice.

EXTERNAL LINK

https://mmpc.org/shared/document.aspx?id=141&docType=Protocol

MATERIALS

NAME CATALOG # VENDOR **CAS NUMBER** RRID NDC0409-7935-19 20% Dextrose injection USP Hospira(Pfizer)

MATERIALS TEXT

Note:

Hospira, RRID:SCR_003985

- Mice may be fasted overnight (~15 hours) or for 5 hours prior to the start of experiment.
- 2 Collect plasma sample (10 µl) before the start of experiment (basal-0 min) to measure basal glucose levels.
- Administer intraperitoneal injection of 20% dextrose (1 or 2 g/kg body weight) using an insulin syringe. 3
- Collect plasma samples (10 µl) at 10, 20, 30, 60, 90, and 120 min following injection to measure circulating glucose concentrations.
- For data analysis, plasma glucose levels vs. time after injection are plotted, and areaunder-curve may be calculated to estimate insulin sensitivity.
- Area-under-curve of glucose tolerance test may be inversely correlated with insulin sensitivity assuming unaffected insulin secretion and pancreatic β-cell function in mice.

This is an open access protocol distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited