

# Separation of free virus particles from sediments in aquatic systems

Roberto Danovaro and Mathias Middelboe

## Abstract

This protocol is for the separation of free virus particles from sediments. It is from:

Danovaro, R., and M. Middelboe. 2010. Separation of free virus particles from sediments in aquatic systems, p. 74–81. In S. W. Wilhelm, M. G. Weinbauer, and C. A. Suttle [eds.], *Manual of Aquatic Viral Ecology*. ASLO.

Please see the [full chapter](#) for additional details.


**Citation:** Roberto Danovaro and Mathias Middelboe Separation of free virus particles from sediments in aquatic systems. **protocols.io**

[dx.doi.org/10.17504/protocols.io.dpg5jv](https://dx.doi.org/10.17504/protocols.io.dpg5jv)

**Published:** 30 Nov 2015

## Guidelines

The optimized protocol for the separation of free virus particles from sediments is shown in Figure 1.



**Figure 1:** Protocol illustrating the steps required for the separation of viruses from the sediment particle and subsequent counting.

## References

- Danovaro, R., A. Dell'Anno, A. Trucco, and S. Vannucci. 2001. Determination of virus abundance in marine sediments. *Appl. Environ. Microbiol.* 67:1384-1387.
- Duhamel, S., and S. Jacquet. 2006. Flow cytometric analysis of bacteria and virus-like particles in lake sediments. *J. Microbiol. Met.* 64:316-322
- Fischer, U. R., A. K. T. Kirschner, and B. Velimirov. 2005. Optimization of extraction and estimation of viruses in silty freshwater sediments. *Aquat. Microb. Ecol.* 40:207-216.
- Hewson, I. and J. A. Fuhrman. 2003. Viriobenthos production and virioplankton sorptive scavenging by suspended sediment particles in coastal and pelagic waters. *Microb. Ecol.* 46:337-347.
- Middelboe, M., R. N. Glud, and K. Finster. 2003. Distribution of viruses and bacteria in relation to diagenetic activity in an estuarine sediment. *Limnol. Oceanogr.* 48:1447-1456.
- Middelboe, M., R. N. Glud 2006. Viral activity along a trophic gradient in continental margin sediments off central Chile. *Mar. Biol. Res.* 2:41-51.

Middelboe, M., R. N. Glud., F. Wenzhöfer, K. Oguri, and H. Kitazato. 2006. Spatial distribution and activity of viruses in the deep-sea sediments of Sagami Bay, Japan. *Deep-Sea Res.* 53:1-13.

Noble, R. T., and J. A. Fuhrman. 1998. Use of SYBR Green I for rapid epifluorescence counts of marine viruses and bacteria. *Aquat. Microb. Ecol.* 14:113-118.

Patel, A., R. T. Noble, J. A. Steele, M. S. Schwalbach, I. Hewson, and J. A. Fuhrman. 2007. Virus and prokaryote enumeration from planktonic aquatic environments by epifluorescence microscopy with SYBR Green I. *Nature Protocols* 2:269-276.

Suttle, C. A. and J. A. Fuhrman. 2010. Enumeration of virus particles in aquatic or sediment samples by epifluorescence microscopy, p. 145-153. In S. W. Wilhelm, M. G. Weinbauer, and C. A. Suttle [eds.], *Manual of Aquatic Viral Ecology*. ASLO.

## Protocol

### Step 1.

Create sediment slurries by combining 0.5 mL sediment and 4.5 mL virus-free water.

### Step 2.

Add tetrasodium pyrophosphate (5 mM final concentration).

### Step 3.

Incubate 15 mins. in the dark.

 DURATION

00:15:00

### Step 4.

Sonicate at 47 KHz for 1 min. (1/3)

 DURATION

00:01:00

### Step 5.

Manually shake for 30 sec.

 DURATION

00:00:30

### Step 6.

Sonicate a second time at 47 KHz for 1 min.

 DURATION

00:01:00

### Step 7.

Manually shake a second time for 30 sec.

 DURATION

00:00:30

### Step 8.

Sonicate a third time at 47 KHz for 1 min.

 DURATION

00:01:00

### Step 9.

Manually shake a third time for 30 sec.

 DURATION

00:00:30

**Step 10.**

Add DNase-I (0.1-0.5 U/mL<sup>-1</sup> final concentration).

**Step 11.**

Incubate 15 mins. in the dark.

 DURATION

00:15:00

**Step 12.**

Dilute supernatant in virus-free seawater (from 10x to 1000x).