

## Species delimitation of Ophiothrix using bPTP 👄

PLOS One

Renata Alitto<sup>1</sup>, Michela Borges<sup>1</sup>, Letícia Dias de Oliveira<sup>1</sup>, Karin Regina Seger<sup>1</sup>, Luciana Bolsoni Lourenço<sup>1</sup>

<sup>1</sup>Universidade Estadual de Campinas

dx.doi.org/10.17504/protocols.io.t2jeqcn



Renata Alitto

#### **ABSTRACT**

This test adds Bayesian support (BS) values to the nodes of the input tree and delimit species based on the Phylogenetic Species Concept. Higher BS values indicate all descendants from the node are more likely to be from one species

**EXTERNAL LINK** 

https://doi.org/10.1371/journal.pone.0210331

THIS PROTOCOL ACCOMPANIES THE FOLLOWING PUBLICATION

Alitto RAdS, Amaral ACZ, Oliveira LDd, Serrano H, Seger KR, Guilherme PDB, Domenico MD, Christensen AB, Lourenço LB, Tavares M, Borges M (2019) Atlantic West Ophiothrix spp. in the scope of integrative taxonomy: Confirming the existence of Ophiothrix trindadensis Tommasi, 1970. PLoS ONE 14(1): e0210331. doi: 10.1371/journal.pone.0210331

PROTOCOL STATUS

### Working

### Phylogenetic tree

The phylogenetic tree generated by the Bayesian Inference was submitted to the Bayesian Poisson Tree Processes - bPTP [56] to test species boundaries.

# Analyses

The analyses were conducted on the web server (available at <a href="http://species.h-its.org/ptp/">http://species.h-its.org/ptp/</a>). The parameters for the run were 500000 MCMC 2 generations, thinning of 100, and Burn-in of 0.25.

This is an open access protocol distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited