



Jun 26,
2019

GG2 - CRISPR-Cas9 episome cloning using red-blue screening for *Phaeodactylum tricornutum*

Mark Moosburner¹, Andrew Allen¹

¹University of California, San Diego

Working

dx.doi.org/10.17504/protocols.io.4acgsaw

JCVI West Protocols



Mark Moosburner

Scripps Institution of Oceanography, J. Craig Venter Institu...



ABSTRACT

To ensure Cas9 activity upon episomal transformation, the 3' end of the Cas9 coding sequencing was transcriptionally fused to an antibiotic selectable marker, shble, by the 2A peptide. The Cas9-2A-shble construct was cloned into a *Phaeodactylum* episome that included a bacterial expression cassette for a red fluorescent protein (RFP). The RFP was also flanked by two BsaI restriction digest sites. The RFP vector with Cas9 was used as a cloning vector to assembly one or more sgRNA *Phaeodactylum* expression cassettes and a single LacZ bacterial expression cassette in place of the RFP. Upon successful assembly, the correct colonies appear blue on selective agar plates while incorrect colonies appear either red or colorless. The golden gate assembly protocol used to synthesize the episome (g51092-2, g24739-A) is as follows:

GG2_ sgRNA cloning into
Cas9 episome using red-
blue screening.pdf



This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited