

The Healthy Brain Network Serial Scanning Initiative, Session 1 Version 2

David O'Connor, Natan Vega Potler, Meagan Kovacs, Ting Xu, Lei Ai, John Pellman, Tamara Vanderwal, Lucas Parra, Samantha Cohen, Satrajit Ghosh, Jasmine Escalera, Natalie Grant-Villegas, Yael Osman, Anastasia Bui, R Cameron Craddock, Michael P Milham

Abstract

This protocol describes MRI and ADHD Quotient Test for Session 1 of the following work:

David O'Connor, et. al. (2017) The Healthy Brain Network Serial Scanning Initiative. *GigaScience*...

Citation: David O'Connor, Natan Vega Potler, Meagan Kovacs, Ting Xu, Lei Ai, John Pellman, Tamara Vanderwal, Lucas Parra, Samantha Cohen, Satrajit Ghosh, Jasmine Escalera, Natalie Grant-Villegas, Yael Osman, Anastasia Bui, R Cameron Craddock, Michael P Milham The Healthy Brain Network Serial Scanning Initiative, Session 1. **protocols.io**

dx.doi.org/10.17504/protocols.io.gxtbxnn

Published: 17 Jan 2017

Before start

Participant offered use of restroom

Protocol

Pre-Scan

Step 1.

Consent is acquired and pregnancy test administered to female participants if they are of childbearing age

⊕ NOTES

David O'Connor 11 Jan 2017

Before Start: Participant offered use of restroom

Pre-Scan

Step 2.

Participant is scanned for magnetic materials, asked to change into scrubs if necessary, and enters the MRI machine

Scan

Step 3.

Localizer



NOTES

David O'Connor 11 Jan 2017

Not Shared, calibration only

Scan

Step 4.

Head Motion training - participant is scanned using an fmri sequence, and presented with visual feedback based on the amount their head moves.



NOTES

David O'Connor 11 Jan 2017

Not Shared, for training purposes only

Scan

Step 5.

Localizer



NOTES

David O'Connor 11 Jan 2017

Not Shared, calibration only

Scan

Step 6.

Resting State - stimuli and scan protocol shared online
(http://fcon_1000.projects.nitrc.org/indi/hbn_ssi/index.html)

Scan

Step 7.

HCP Task - participant was scanned with fmri sequence while performing one of seven tasks used in the Human Connectome project
(<http://www.humanconnectome.org/documentation/Q1/task-fMRI-protocol-details.html>)

Scan

Step 8.

B0 field map

Scan

Step 9.

T1 weighted Multiecho MPRAGE - scan protocol shared online
(http://fcon_1000.projects.nitrc.org/indi/hbn_ssi/index.html)

Scan

Step 10.

DWI- scan protocol shared online (http://fcon_1000.projects.nitrc.org/indi/hbn_ssi/index.html)

Scan

Step 11.

DKI- scan protocol shared online (http://fcon_1000.projects.nitrc.org/indi/hbn_ssi/index.html)

Scan

Step 12.

Whole brain T1 - scan protocol shared online
(http://fcon_1000.projects.nitrc.org/indi/hbn_ssi/index.html)

Scan

Step 13.

Inversion Recovery - scan protocol shared online
(http://fcon_1000.projects.nitrc.org/indi/hbn_ssi/index.html)

Scan

Step 14.

Whole brain T2 - scan protocol shared online
(http://fcon_1000.projects.nitrc.org/indi/hbn_ssi/index.html)

Scan

Step 15.

Magnetization transfer (OFF) - - scan protocol shared online
(http://fcon_1000.projects.nitrc.org/indi/hbn_ssi/index.html)

Scan

Step 16.

Magnetization transfer (ON) - - scan protocol shared online
(http://fcon_1000.projects.nitrc.org/indi/hbn_ssi/index.html)

Scan

Step 17.

T2 FLAIR - scan protocol shared online (http://fcon_1000.projects.nitrc.org/indi/hbn_ssi/index.html)

Post-Scan

Step 18.

Removed from Scanner

Post-Scan

Step 19.

Quotient - Quotient is a computer based task designed to assess three core symptoms of ADHD: hyperactivity, attention and impulsivity. Participants respond to stimuli presented with random timing and random placement on a screen. Completion of the task takes up to 30 minutes.

🔗 NOTES

David O'Connor 11 Jan 2017

DUA required for this data