



May 10,
2019

Working

U Mass - Insulin clearance [↗](#)

Jason Kim¹

¹University of Massachusetts

dx.doi.org/10.17504/protocols.io.x2wfqfe

Mouse Metabolic Phenotyping Centers

Tech. support email: info@mmpc.org

Lili Liang

ABSTRACT

Summary:

Insulin clearance test measure systemic clearance of insulin following a bolus injection. Liver accounts for the majority of systemic insulin clearance following secretion from pancreatic β - cells into portal circulation. Hepatic clearance of insulin may be affected by obesity and in other mouse models of altered metabolism. Alterations in insulin clearance may also affect glucose and lipid metabolism.

EXTERNAL LINK

<http://mmpc.org/shared/document.aspx?id=140&docType=Protocol>

MATERIALS

NAME	CATALOG #	VENDOR	CAS NUMBER	RRID
Insulin	Regular human insulin, U-100	Novolin		
Insulin Ultrasensitive ELIZA	80-INSMSU-E01	Alpco		AB_2792981

MATERIALS TEXT

Note:

Cite this, **Alpco Diagnostics Cat# 80-INSMSU-E01**, [RRID:AB_2792981](#)

- 1 Mice are fasted for 5 hours prior to the start of experiment.
- 2 Collect plasma sample (20 μ l) before the start of experiment (basal-0 min) to measure basal insulin and glucose levels.
- 3 Administer intraperitoneal injection of insulin (0.5 or 0.75 unit/kg body weight) using an insulin syringe.
- 4 Collect plasma samples (10 μ l) at 10, 20, 30, 60, 90 and 120 min for the measurement of plasma insulin concentrations.
- 5 For data analysis, plasma insulin levels vs. time after insulin injection are plotted, and area-under-curve may be calculated to estimate systemic insulin clearance.



This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits

