



Mar 07, 2019

Working

RNA extraction with PGTX

Lutz Berwanger¹

¹Heinrich-Heine Universität Düsseldorf

dx.doi.org/10.17504/protocols.io.ywwfxfe

M4454 - Synthetische Biologie und Biotechnologie



Sebastian Triesch Institute for Synthetic Microbiology, HHU Düsseldorf





ABSTRACT

This protocol is used for RNA extraction in cyanobacteria after Pinto et al. 2009.

Pinto, Fernando Lopes; Thapper, Anders; Sontheim, Wolfgang; Lindblad, Peter (2009): Analysis of current and alternative phenol based RNA extraction methodologies for cyanobacteria. In: BMC molecular biology 10, S. 79. DOI: 10.1186/1471-2199-10-79



PROTOCOL STATUS

Working

We use this protocol in our group and it is working

STEPS MATERIALS

NAME CATALOG # **VENDOR**

Chloroform

Isopropanol

Ethyl alcohol, Pure 200 proof, for molecular biology E7023 Sigma Aldrich

SAFETY WARNINGS

Additional Information

Always provide RNA work on ice and always wear Gloves.









GHS Label elements, including precautionary statements

Signal word

Danger



Hazard statement(s)

H227 Combustible liquid.
H302 Harmful if swallowed.
H319 Causes serious eye irritation.
H311 Toxic in contact with skin.

H314 Causes severe skin burns and eye damage.

H330 Fatal if inhaled.

H341 Suspected of causing genetic defects.

H371 May cause damage to organs.

H373 May cause damage to organs through prolonged or repeated exposure.

H402 Harmful to aquatic life.

Precaution Statement(s)

P260 Do not breathe dust/ fume/ gas/ mist/ vapours/ spray.

P280 Wear protective gloves/ protective clothing/ eye protection/ face protection.

P284 Wear respiratory protection.

P305 + P351 + P338 IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do.

Continue rinsing.

P310 Immediately call a POISON CENTER or doctor/ physician.

Material list

Chemical	Volume/Mass	Company	Serial no.	Comments
Phenol	39.6g			See add. Info
Glycerol	6.9mL			
8-hydroxyquinoline	0.1g			
EDTA	0.58g			See add. Info
Sodium acetate	0.8g			
Guanidine thiocyanate	9.5g			
Guanidine	4.6g			
hydrochloride				
Chloroform				See add. Info
Isopropanol				See add. Info
EtOH	70%			
DEPC treated Water		Roth		

Buffer

Name	Ingredients		Comments
100 mL PGTX Solution:	Phenol	39.6 g	
	Glycerol	6.9 mL	
	8-hydroxyquinoline	0.1 g	
	EDTA	0.58 g	
	Sodium acetate	0.8 g	
	Guanidine thiocyanate	9.5 g	
	Guanidine hydrochloride	4.6 g	

Equipment list

Device	Company	Comments
Centrifuge	Eppendorf	4°C
Thermo Mix	Eppendorf	95°C
Centrifuge	Heraeus	4°C
Freezer	Liebherr	-20°C

Fill 50 mL Falcon tube with ice. Fill with bacterial liquid culture, up to a volume of ~45 mL. Centrifuge for 3 minutes at 4°C and 4.000 g. **© 00:03:00** Discard supernatant. Resuspend cell pellet in residual water (~1 mL). Transfer to 2 mL 'safe lock' tube. 4 **© 00:00:15** Resuspend the cell pellet in 1 ml of PGTX. Freeze in liquid nitrogen and store at -20 °C. **▲**SAFETY INFORMATION Wear goggles, a lab coat and gloves when dealing with PGTX and liquid nitrogen. Incubate for 5 min at 95 °C, shaking at 250 rpm in Thermomixer (Eppendorf) **© 00:05:00** Rapidly chill 5 min on ice. **© 00:05:00** Add 700 µl Chloroform. **A**SAFETY INFORMATION Wear goggles, a lab coat and gloves when dealing with phenol and chloroform. Chloroform Let the samples incubate for 10 min at room temperature in a Thermomixer. Vortex from time to time. **© 00:10:00** $Centrifuge \ for \ 15 \ min \ at \ 14.000 \ g, \ 4 \ ^\circ C. \ Transfer \ the \ upper \ aqueous \ phase \ (\textbf{contains RNA}) \ to \ a \ fresh \ reaction \ tube \ and \ add \ the \ same$ 10 volume (450 μ L) of Aqua- P/C/I (Phenol/Chloroform/Isoamylalcohol). **© 00:15:00 ▲**SAFETY INFORMATION Wear goggles, a lab coat and gloves when dealing with phenol and chloroform. Thoroughly mix by vortexing. Centrifuge for 15 min at high speed. Transfer the upper aqueous phase to a 1.5 mL reaction tube. 11 **© 00:15:00 ▲**SAFETY INFORMATION Wear goggles, a lab coat and gloves when dealing with phenol and chloroform. 12 Add 1 volume of isopropanol.

03/07/2019

✓ protocols.io



13 Mix and incubate for at least 30 min at -20 °C. (Can be left overnight.)

© 00:30:00

14 Centrifuge at 14.000 g for 30 min.

© 00:30:00

- 15 Discard supernatant.
- 16 Wash with 75% chilled ethanol. Avoid resuspending the pellet.



17 Centrifuge at 14.000 g.

© 00:05:00

18 Repeat washing step with 75% chilled ethanol.



19 Air dry pellet at RT. Do not overdry!

© 00:10:00

20 $\,\,$ Resuspend the pellet with 30 μL volume of ddH20.



This is an open access protocol distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited