



Working

Dengue genotyping by E gene amplification and sequencing [↗](#)

PLOS Neglected Tropical Diseases

Jeyanthi Suppiah¹

¹Institute for Medical Research, Kuala Lumpur; Universiti Putra Malaysia, Selangor.

Sep 19, 2018 [dx.doi.org/10.17504/protocols.io.racd2aw](https://doi.org/10.17504/protocols.io.racd2aw)



Jeyanthi Suppiah

ABSTRACT

- 1) The partial E gene of dengue virus is amplified before sequencing by using four sets of serotype-specific oligonucleotides as referenced in the manuscript text.
 - 2) The reaction mixture is prepared as the following:
 - 12.5ul of MyTaq RT-PCR (Bioline, Korea)
 - 4.5ul of Nuclease-Free Water
 - 1.0 ul of Forward Primer for the respective serotype (10uM)
 - 1.0 ul of Reverse Primer for the respective serotype (10uM)
 - 0.5ul of RNase inhibitor
 - 0.5ul of Reverse Transcriptase (RT)
 - 3) Aliquot 20ul in each PCR tubes.
 - 4) 5ul of Dengue RNA (with known serotype) is added to the corresponding PCR tube.
 - 5) PCR amplification is performed on CFX-96 (BioRad) conventional PCR thermal cycler.
 - 6) The PCR is performed with the following cycling profile: 35 cycles (94 °C for 1 min, 55 °C for 1 min and 72 °C for 1.5 min) followed by an extension reaction at 72 °C for 5 min.
 - 7) A 25 µl aliquot of each PCR reaction is analyzed on 1.5 % pre-stained agarose by gel electrophoresis, run for 30 min at 90 V.
 - 8) The gel is then viewed under UV illumination.
 - 9) The expected amplicon size for each serotype is 578bp for DENV1, 617bp for DENV 2, 582bp for DENV3 and 572bp for DENV 4
- The corresponding amplicons are extracted from the agarose gel and purified by Gel Extraction Kit (Qiagen, USA) according to the manufacturer's instruction.
- 10) The final elution volume is 30 µl of purified PCR amplicons. Then, 5 µl of these are reanalyzed on 1.5 % agarose gel to substantiate the accuracy of purification step. The purified PCR amplicons were outsourced for sequencing.

EXTERNAL LINK

<https://doi.org/10.1371/journal.pntd.0006817>

THIS PROTOCOL ACCOMPANIES THE FOLLOWING PUBLICATION

Suppiah J, Ching S, Amin-Nordin S, Mat-Nor L, Ahmad-Najimudin N, Low GK, Abdul-Wahid M, Thayan R, Chee H (2018) Clinical manifestations of dengue in relation to dengue serotype and genotype in Malaysia: A retrospective observational study. PLoS Negl Trop Dis 12(9): e0006817. doi: [10.1371/journal.pntd.0006817](https://doi.org/10.1371/journal.pntd.0006817)

PROTOCOL STATUS

Working



This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited