

# Confronting data with reality: Spatial patterning in simulated ridge and slough landscape from RASCAL everglades modeling

Jenna T. Baughman and Laurel G. Larsen

DS421 - Summer 2016 RET

## Introduction

The Florida Everglades have been described as having regular or periodic landscape patterning, yet, the processes that develop and maintain it are not well understood. Quantifying the spatial characteristics of these patterns can provide insight into the processes and provide metrics for model validation. Here we used spatial spectral analyses to compare landscape patterns and frequencies represented in simulated ridges and sloughs from the RASCAL Everglades model to observations from areal imagery.



Figure 1: Google Earth aerial image demonstrating a ridge and slough landscape from the Everglades Wildlife Management Area in southern Florida. Darker colors are ridges dominated by sedge sawgrass (*Cladium jamaicense*) and lighter colors are sloughs, containing less dense emergent and floating macrophytes. Brighter green spots are tree islands.

## Simulations

Everglades ridge and slough landscapes were simulated using the RASCAL model (Larsen and Harvey, 2011).




Figure 2: Sample RASCAL model output and aerial image displaying similar ridge and slough landscapes (Larsen and Harvey, 2011).

## Methods

All analyses were performed using Matlab version R2016A under an academic license.

## Results

### Image Processing




Figure 5: RASCAL model image output (left) and with ridges outlined (right) for perimeter calculations.




Figure 3: Casey et al. (in review) empirical sites converted to binary images.




Figure 4: Patch (ridge) area complementary cumulative distribution function (CCDF).

## Research Questions:

1. Is there evidence of **regular patterning** in RASCAL everglades landscape images?
2. Is there a **dominant spatial frequency** in RASCAL everglades landscape images?
3. How do these results **compare** to the same analyses on aerial images or vegetation maps?
4. How do these results vary with **different RASCAL parameters**?




Figure 6: Casey et al. (in review) perimeter to area scaling indicates ridge perimeters increase faster than expected for fractal relationship.




Figure 7: Casey et al. (in review) omni-directional  $r$  spectra show no support for periodic behavior (indicated by absence of strong peaks).

### Area CCDF




Figure 9: Ridge sizes show strong support for both the generalized Pareto distribution (shown in red) and a lognormal distribution (shown in blue).

### Spatial Spectral Analyses




Figure 10: Omni-directional  $r$  spectrum indicates presence of a strong peak and support for periodic behavior.

### Perimeter to Area Scaling




Figure 8: Perimeter to area scaling shows a linear relationship (linear model is shown in blue; quadratic model in red) indicating that ridge perimeter does not increase differently than expected for a fractal relationship.

## Next Steps

- Analyze **ridge elongation** statistics
- Compute **directional  $r$  spectrum**; perpendicular to flow and find best **fit** for data
- Repeat all analyses with different RASCAL images resulting from **different input parameters**
- **Interpret** results!

## References

Casey, S.T., Cohen, M.J., Acharya, S., Kaplan, D.A., and Jawitz, J.W., in review. *Hydrology and Earth System Sciences*.  
Larsen, L.G., and Harvey, J.W., 2011. *Geomorphology* 126:279-296.