



Linear Regression

NYU K12 STEM Education: Machine Learning

Department of Electrical and Computer Engineering,
NYU Tandon School of Engineering
Brooklyn, New York

- ▶ [Course Website](#)
- ▶ Instructors:



Rugved Mhatre

rugved.mhatre@nyu.edu



Akshath Mahajan

avm6288@nyu.edu

Outline

1. Python

2. Statistics

3. Linear Regression

4. Multivariable Linear Regression

Vectorize Programming

Demo on Vectorize Programming

Plotting Functions

Generate and plot the following functions in Python:

- ▶ Scatter plot of points: $(0, 1), (2, 3), (5, 2), (4, 1)$
- ▶ Straight Line: $y = mx + b$
- ▶ Sine-wave: $y = \sin(x)$
- ▶ Polynomial e.g. $y = x^3 + 2$
- ▶ Exponential e.g. $y = e^{-2x}$
- ▶ Choose a function of your own

Use Wikipedia and NumPy documentation to search for mathematical formulas and python functions

Outline

1. Python

2. Statistics

3. Linear Regression

4. Multivariable Linear Regression

Mean

The mean, or average, is the sum of all the values in a dataset divided by the number of values. It provides a measure of the central tendency of the data.

$$\text{Mean}(\mu) = \frac{1}{N} \sum_{i=1}^N x_i$$

Example:

For the dataset $X = [2, 4, 6, 8, 10]$,

$$\mu = \frac{2 + 4 + 6 + 8 + 10}{5} = 6$$

Variance

Variance measures the spread of the data points around the mean. It indicates how much the data varies from the mean.


$$\text{Variance}(\sigma^2) = \frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2$$

Example:


For the dataset $X = [2, 4, 6, 8, 10]$,

$$\sigma^2 = \frac{(2 - 6)^2 + (4 - 6)^2 + (6 - 6)^2 + (8 - 6)^2 + (10 - 6)^2}{5} = 8$$


Mean and Variance Visualization



Mean and Variance Visualization



Mean and Variance Visualization



Standard Deviation

Standard deviation is the square root of the variance. It provides a measure of the spread of the data points in the same units as the data itself.


$$\text{Standard Deviation}(\sigma) = \sqrt{\text{Variance}}$$

Example:


Using the variance calculated previously,

$$\sigma = \sqrt{8} \approx 2.83$$

Standard Deviation Visualization



Standard Deviation Visualization



Covariance

Covariance measures the degree to which two variables change together. If the covariance is positive, the variables tend to increase together; if negative, one variable tends to increase when the other decreases.




$$\text{Covariance}(\text{Cov}(X, Y)) = \frac{1}{N} \sum_{i=1}^N (x_i - \mu_X)(y_i - \mu_Y)$$

Example:

For the datasets $X = [2, 4, 6]$ and $Y = [3, 6, 9]$,

$$\text{Cov}(X, Y) = \frac{(2 - 4)(3 - 6) + (4 - 4)(6 - 6) + (6 - 4)(9 - 6)}{3} = 6$$

Covariance Visualization



Looking at our ice-breaker data in spreadsheets

- ▶ Columns have labels in the first row
- ▶ Collected data (numbers, words) follow below
- ▶ Let's export it to a Comma-Separated Values (CSV) file and open it

Outline

1. Python

2. Statistics

3. Linear Regression

4. Multivariable Linear Regression

Linear Regression in a nutshell

- ▶ Consider a function $y = 2x + 1$
- ▶ Here we introduce a new notation $f(x) = 2x + 1$
- ▶ What this means is that we have a function $f(x)$ which has x as its variable.
- ▶ If we have different x values we will have different values of $f(x)$

Linear Regression in a nutshell

- ▶ For $f(x) = 2x + 1$ and setting $x = 1$ we have $f(x) = 3$
- ▶ For $f(x) = 2x + 1$ and setting $x = 0$ we have $f(x) = 1$
- ▶ For $f(x) = 2x + 1$ and setting $x = -1.5$ we have $f(x) = -2$

Linear Regression in a nutshell

- ▶ We believe that dataset are representation of underlying models which can be represented as functions of features.
- ▶ For example, we can build a model to forecast weather, we can use the features humidity, current temperature and wind speed to estimate what the temperature will be tomorrow.
- ▶ Here we have $f(x)$ representing the tomorrow's temperature and x being a vector containing humidity, current temperature and wind speed.

Linear Regression in a nutshell

- ▶ But many times we do have $f(x)$ available, our task here is to figure out what $f(x)$ is using the data available to us.
- ▶ Here $f(x)$ is called a model.
- ▶ In other words, we want to find a model that fits the data.

Linear Regression in a nutshell

- ▶ It would be easier to have a “framework” of the model ready and find the model parameters using the data.
 - $f(x) = w_1x + w_0$
 - $f(x) = w_2x^2 + w_1x + w_0$
 - $f(x) = \frac{1}{e^{-(w_1x+w_0)}+1}$
- ▶ The numbers w_0 , w_1 and w_2 are called model parameters.
- ▶ We often write the model as $f(x; w)$, stacking all parameters to a vector w .

Structure of a dataset

- ▶ In a dataset we have many data.
- ▶ We can represent each piece of data as (x_i, y_i) , $i = 1, 2, 3, \dots$
- ▶ x_i is called the feature and y_i is called the label.
- ▶ The relationship between x_i and y_i and the model f is $f(x_i) = y_i$
- ▶ For example, if the weather forecast says it will be 21°C (69.8°F) if it turns out to be 22°C (71.6°F) you won't be yelling at the TV.

How would you fit a line?

Can you find a line that passes through $(0, 0)$ and $(1, 1)$?

- ▶ The “framework” of the model is $f(x) = w_1x + w_0$
- ▶ The data is $(x = 0, f(x) = y = 0)$ and $(x = 1, f(x) = y = 1)$.
- ▶ The process of finding a model to fit the data is to find the values of w_1 and w_0 .

How would you fit a quadratic curve?

Can you find a quadratic curve that passes through $(0, 0)$, $(1, 1)$ and $(-1, 1)$?

- ▶ The “framework” of the model is $f(x) = w_2x^2 + w_1x + w_0$
- ▶ The data is $(x = 0, f(x) = y = 0)$, $(x = 1, f(x) = y = 1)$ and $(x = -1, f(x) = y = 1)$
- ▶ The process of finding a model to fit the data is to find the values of w_2 , w_1 and w_0 .

What model do we use for this dataset?

- ▶ Open [Linear Regression Demo](#)
- ▶ Can you find a line that goes through ALL of the data points? Why?

Is Your Model a Good Fit?

- ▶ How would you determine if your model is a good fit or not?
 - How will you determine this?
 - Is there a quantitative way?
- ▶ We now introduce a new notation $f(x_i) = \hat{y}_i$ here the $\hat{\cdot}$ represents $f(x_i)$ is a prediction of y_i .

Error Functions

- ▶ An **error function** quantifies the discrepancy between your model and the data.
 - They are non-negative, and go to zero as the model gets better.
- ▶ Common Error Functions:
 - Mean Squared Error:

$$\text{MSE} = \frac{1}{N} \sum_{i=1}^N ||y_i - \hat{y}_i||^2$$

- Mean Absolute Error:

$$\text{MAE} = \frac{1}{N} \sum_{i=1}^N |y_i - \hat{y}_i|$$

- ▶ In later units, we will refer to these as **cost functions** or **loss functions**.

Linear Regression

- ▶ Linear models: For scalar-valued feature x , this is $f(x) = w_1x + w_0$
- ▶ One of the simplest machine learning model, yet very powerful.

Least Square Solution

- Model:

$$f(x) = w_1x + w_0$$


- Loss:

$$J(w_0, w_1) = \frac{1}{N} \sum_{i=1}^N ||y_i - f(x_i)||^2$$

- Optimization: Find w_0, w_1 such that $J(w_0, w_1)$ is the least possible value (hence the name “least square”).

Loss Landscape

$$J(w_0, w_1) = \frac{1}{N} \sum_{i=1}^N \|y_i - f(x_i)\|^2$$



Least Square Solution: Using Pseudo-Inverse

For N data points (x_i, y_i) we have,

$$\hat{y}_1 = w_0 + w_1 x_1$$

$$\hat{y}_2 = w_0 + w_1 x_2$$

$$\vdots$$

$$\hat{y}_N = w_0 + w_1 x_N$$

Least Square Solution: Using Pseudo-Inverse

In matrix form we have,

$$\begin{bmatrix} \hat{y}_1 \\ \hat{y}_2 \\ \vdots \\ \hat{y}_N \end{bmatrix} = \begin{bmatrix} 1 & x_1 \\ 1 & x_2 \\ \vdots & \vdots \\ 1 & x_N \end{bmatrix} \begin{bmatrix} w_0 \\ w_1 \end{bmatrix}$$

We can write it as $\hat{Y} = X \times W$. We call X the design matrix.

Least Square Solution: Using Pseudo-Inverse

- ▶ We can put the desired labels in matrix form as well:

$$Y = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_N \end{bmatrix}$$

- ▶ Our goal is to minimize the error between Y and \hat{Y} which can be written as $\|Y - \hat{Y}\|^2$
- ▶ **Exercise:** Verify

$$\|Y - \hat{Y}\|^2 = \|Y - XW\|^2 = \sum_{i=1}^N \|y_i - (w_0 + w_1 x_i)\|^2$$

Linear Least Square

$$\min_W \frac{1}{N} \|Y - XW\|^2$$

The solution looks like this,

$$W = (X^T X)^{-1} X^T Y$$

Outline

1. Python
2. Statistics
3. Linear Regression
4. Multivariable Linear Regression

Multivariable Linear Regression

What if we have multivariable data with x being a vector?

Example:

$$x_i = \begin{bmatrix} x_{i1} \\ x_{i2} \end{bmatrix}$$

$$\hat{y}_1 = w_0 + w_1x_{11} + w_2x_{12}$$

$$\hat{y}_2 = w_0 + w_1x_{21} + w_2x_{22}$$

$$\vdots$$

$$\hat{y}_N = w_0 + w_1x_{N1} + w_2x_{N2}$$

Multivariable Linear Regression

The model can be written in matrix-vector form as:

$$\begin{bmatrix} \hat{y}_1 \\ \hat{y}_2 \\ \vdots \\ \hat{y}_N \end{bmatrix} = \begin{bmatrix} 1 & x_{11} & x_{12} \\ 1 & x_{21} & x_{22} \\ \vdots & \vdots & \vdots \\ 1 & x_{N1} & x_{N2} \end{bmatrix} \begin{bmatrix} w_0 \\ w_1 \\ w_2 \end{bmatrix}$$

Solution remains the same $W = (X^T X)^{-1} X^T Y$

► **Exercise:** Open [Multivariable Linear Regression Demo](#)